

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
3 January 2002 (03.01.2002)

PCT

(10) International Publication Number
WO 02/00924 A2

- (51) International Patent Classification⁷: C12Q 1/68 (74) Agent: MBM & CO.; P.O. Box 809, Station B, Ottawa, Ontario K1P 5P9 (CA).
- (21) International Application Number: PCT/CA01/00932
- (22) International Filing Date: 28 June 2001 (28.06.2001)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
2,310,263 28 June 2000 (28.06.2000) CA
- (71) Applicant (for all designated States except US): INSTITUT NATIONAL DE LA RECHERCHE SCIENTIFIQUE [CA/CA]; Tour de la Cité, 2600 Boulevard Laurier, Bureau 640, Case Postale 7500, Sainte-Foy, Québec G1V 4C7 (CA).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): TIJSEN, Peter [NL/CA]; 76 Winston Circle, Pointe-Claire, Québec H9S 4X6 (CA). ZADORI, Zoltan [HU/CA]; 3215 Boulevard Cartier Ouest, App. 106, Laval, Québec H7V 1J8 (CA).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:
— without international search report and to be republished upon receipt of that report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 02/00924 A2

(54) Title: VIRAL PHOSPHOLIPASE A₂ ENZYMES, ANTI-VIRAL AGENTS AND METHODS OF USE

(57) Abstract: The present invention provides a novel class of phospholipase A₂ enzyme (PLA₂) from viruses and a method for identifying, isolating, purifying and characterizing enzymes of this class. The present invention includes viral PLA₂ proteins, nucleic acids, and antisense oligonucleotides, and the use of these molecules in screening methods for anti-viral agents, in decreasing the infectivity and/or replication of viruses and as research tools. The present invention further includes treatment or prevention of virus-associated diseases using viral PLA₂ inhibitors and the use of the viral PLA₂-encoding region to improve virus-based gene therapy vectors.

VIRAL PHOSPHOLIPASE A₂ ENZYMES, ANTI-VIRAL AGENTS AND METHODS OF USE

FIELD OF THE INVENTION

The present invention pertains to a class of phospholipase A₂ proteins and nucleic acids and
5 their use. More particularly the present invention pertains to viral phospholipase A₂ proteins
and nucleic acids and their use.

BACKGROUND OF THE INVENTION

Phospholipase A₂

Phospholipase enzymes catalyze the removal of fatty acid residues from phosphoglycerides.

- 10 Phospholipases A₂ (PLA₂s) are a superfamily of key enzymes involved in a multitude of
(patho)physiological and cellular processes, including lipid membrane metabolism, signal
transduction pathways, inflammation, acute hypersensitivity, and degenerative diseases
(Balsinde *et al.*, (1999) *Annu. Rev. Pharmacol. Toxicol.* **39**:175-189; Kramer and Sharp
(1997) *FEBS Letters* **410**:49-53; Nishizuka (1992) *Science* **258**:607-614; Dennis (1994) *J.*
15 *Biol. Chem.* **269**:13057-13060; Dennis (1997) *Trends Biochem. Sci.* **22**:1-2; Yuan and Tsai
(1999) *Biochim Biophys Acta* **1441**:215).

- PLA₂s hydrolyze the ester bonds at the *sn*-2 position of the glycerol moiety of membrane
phospholipids to yield lysophospholipids and fatty acids, from which secondary messengers
may be generated. These secondary messengers may modulate enzyme activities, ion
20 channels, and (post-)transcription factors (Nishizuka (1992) *Science* **258**:607-614; Rao *et al.*,
(1993) *Oncogenes* **8**:2759-2764; Yao *et al.*, (1995) *Nature* **378**:307-310).

- PLA₂s are found in many living species and form a diverse family of enzymes. The main
groups of PLA₂s are the cytosolic PLA₂s, the Ca²⁺-independent PLA₂s, and the secretory
25 PLA₂s (sPLA₂s). A large number of sPLA₂s, from host cells as well as from snake and bee
venoms, have been structurally characterized, and show a high degree of sequence homology
(See, for example, Chang *et al.*, (1987) *Biochemical Pharmacology* **36**:2429-2436). Although

the known sPLA₂s were originally divided into groups by source organism and their primary amino acid sequences, they are now characterized by a growing list of other attributes, such as the number and position of disulfide bonds. Group I includes pancreatic sPLA₂s from vertebrates, including mammals, and snake venom sPLA₂s. These sPLA₂s are active at pH 5 6-8 and their activity is dependent on the presence of calcium ion (Ca²⁺). Group II contains a mixture of non-pancreatic, or synovial, sPLA₂s and sPLA₂s from the venom of crotalids and viperids. Both group I and group II members have a similar 3D-structure. Group III contains sPLA₂s from bee venoms. sPLA₂s from all three groups have a 14 kDa molecular mass and are disulfide-rich (Dennis (1994) *J. Biol. Chem.* 269:13057-13060; Dennis (1997) *Trends* 10 *Biochem. Sci.* 22:1-2). While all low-molecular weight (~13.5-16.8 kDa), Ca²⁺-dependent PLA₂s possess secretion sequences and have been found extracellularly, some are also cell-associated (Anderson *et al.* (1994) *Prostaglandins, Leukotrienes and Essent Fatty Acids* 51:19; Fayard *et al.* (1998) *J Cell Sci* 111: 985). Other types of PLA₂s include group V PLA₂ (Chen *et al.* (1994) *J. Biol. Chem.* 269:2365), and group X PLA₂ isoforms (Valentin *et al.* 15 (1999) *J. Biol. Chem.* 274: 31195).

The product of sPLA₂ activity, arachadonic acid, is processed into bioactive lipid mediators or shuttled into pathways for the synthesis of eicosanoids. In mammals eicosanoids, such as prostaglandins, prostacyclins, thromboxanes and leukotrienes, are involved in pain, inflammation and fever. As such, inhibitors of mammalian sPLA₂ have been described for 20 use in the treatment of a variety of inflammatory disorders (for example, see International Patent Application No. WO 96/27604, European Patent Application No. EP 950 657 A2 and U.S. Patent No. 5,948,626). In addition, a recent report has shown that sPLA₂ isolated from bee or snake venom inhibits human immunodeficiency virus (HIV) by blocking viral entry into host cells (Fenard *et al.*, (1999) *J. Clin. Invest.* 104:611).

25 While prokaryotic versions of PLA₂ have been reported in bacteria such as *Escherichia coli* and *Streptomyces violaceoruber*, viral PLA₂ has not previously been identified.

Viruses

Viruses are infectious agents that are found in virtually all life forms, including humans,

animals, plants, fungi, and bacteria. Viruses often damage or kill the cells that they infect, causing disease in infected organisms. The difficulty in developing anti-viral therapies stems from the large number of variant viruses that can cause the same disease, as well as the inability of drugs to disable a virus without disabling healthy cells. The development of
5 specific anti-viral agents, therefore, is a major focus of current research.

Therapeutic Anti-Viral Treatments

Although a number of methods exist currently for the treatment of viral infections, many of these are complicated by the side effects the anti-viral agent(s) has on the host system.

One example of a method for inhibiting a viral infection is provided by U.S. Patent No.
10 5,449,608, which describes a method of inhibiting the human parvovirus, B19. The method comprises administering compositions containing a B19 binding domain, thus preventing binding of B19 to host cells. Unfortunately, such an agent competes with similar molecules already present in the host cells, without a clear competitive edge, and is prone to catalytic breakdown.

15 Similarly, a soluble form of the receptor (CD-4) for the human immunodeficiency virus, HIV-1, has been shown to be somewhat effective in reducing the infectivity of HIV in tissue culture, but this result could not be replicated *in vivo*. This is likely due to the large number of receptor binding sites present on a single virus particle, thus making it difficult for a soluble receptor to block all sites and prevent binding of the virus to cells (see Flint *et al.*,
20 *Principles of Virology* (Washington, DC: ASM Press, 1999) 119).

Clearly, a need remains to provide easily applicable methods and agents that may be used to effectively treat viral infections, particularly in cases where vaccination has failed or where subjects were not vaccinated.

The present invention provides a means of inhibiting virus infection and/or replication in
25 animals, including humans, and of treating persistent infections and associated syndromes.

This background information is provided for the purpose of making known information believed by the applicant to be of possible relevance to the present invention. No admission is necessarily intended, nor should be construed, that any of the preceding information constitutes prior art against the present invention.

5 SUMMARY OF THE INVENTION

An object of the present invention is to provide viral phospholipase A₂ enzymes, antiviral agents and methods of use. In accordance with an aspect of the present invention, there is provided a viral polypeptide that has phospholipase A₂ activity.

In accordance with another aspect of the present invention, there is provided an isolated
10 polynucleotide encoding a viral polypeptide that has phospholipase A₂ activity, antisense oligonucleotides complementary to the polynucleotides, vectors comprising the polynucleotides, and host cells genetically engineered with the polynucleotides or vectors.

In accordance with another aspect of the present invention there is provided methods of decreasing parvovirus infection and replication, and thereby methods of treating parvovirus-
15 related diseases.

In accordance with another aspect of the present invention viral PLA₂s are used in manufacturing, clinical, and research settings. At present, bee venom is commonly used as a standard in measuring PLA₂ activity. The viral PLA₂ of porcine parvovirus (PPV), however, has a 40 to 100-fold higher specific activity than bee venom. This high activity
20 makes PPV PLA₂ ideal for use as a PLA₂ standard.

In accordance with another aspect of the present invention there is provided methods for screening compounds in order to identify inhibitors of viral PLA₂ activity, comprising adding a candidate inhibitor compound to a solution containing a viral PLA₂ and a phospholipid substrate, and detecting the inhibition of PLA₂ activity.

In accordance with another aspect of the present invention there is provided antisense oligonucleotides designed to inhibit expression of viral PLA₂.

In accordance with a further aspect of the present invention is the use of these viral PLA₂ inhibiting agents as anti-viral agents in the treatment of various viral infections.

- 5 In accordance with still another aspect of the present invention there is provided the use of viral PLA₂ nucleic acids, or variations thereof, to improve gene-therapy vectors.

Various other objects and advantages of the present invention will become apparent from the detailed description of the invention.

BRIEF DESCRIPTION OF THE DRAWINGS

- 10 **Figure 1** depicts a parvovirus genome.

The right-half of the parvovirus genome contains a large ORF coding for a nested set of capsid proteins (VPs) from alternative in-frame initiation codons so that these VPs differ by their *N*-terminal extensions. The VPs can be subdivided into 3 domains: common *C*-terminal domains required for capsid formation (60 copies per capsid); a small hinge-(like) domain;

- 15 and unique *N*-terminal extensions. The capsid-forming domain consists of a β -barrel in which β -strands (black boxes: αA and αB are helices) are connected by loops (L1-4). About 100 amino acids upstream of the start of VP is a conserved domain (cd). The shaded boxes represent the domains (V1ups) cloned in expression vectors (VP1up: PPV-VP1up amino acids 2-174 [SEQ ID NO: 50]; B19-VP1up amino acids 2-240 [SEQ ID NO: 52]; and
- 20 GmDENV-VP1up amino acids 1-378 [SEQ ID NO: 54]).

Figure 2 presents a protein sequence alignment of parvovirus VP1up regions and representatives of sPLA₂ groups. Parvoviral and sPLA₂ sequences are separated by numbers of the common structural numbering for group I/II sPLA₂ (Renetseder *et al.*, (1985) *Annu. Rev. Pharmacol. Toxicol.* 39:175; Kramer and Sharp (1997) *FEBS Lett.* 410:49; Nishizuka

- 25 (1992) *Science* 258:607). Black shading indicates 100% identity among sPLA₂s. Dark grey

indicates 100% identity among vPLA₂s. Medium grey shading indicates 100% identity among all PLA₂s. GenBank accession numbers are indicated except for those not yet deposited.

Figure 3 demonstrates the specificity of PLA₂ activity.

5 **A.** Thin layer chromatography after hydrolysis of phosphatidylcholine substrate (PL) into fatty acid (FA) and lysophosphatidylcholine (LPL) by expressed pvPLA₂ in the mixed micelles assay. PLA₂s used were: lanes 1, 2 and 5 (0.6 ng of PPV PLA₂), lanes 3 and 4 (250 ng of B19 PLA₂), lane 6 (350 ng of *GmDNV* PLA₂), lane 7 (2,000 ng thioredoxin as negative control) and lane 8 (15 ng bee venom PLA₂). The 3C9 monoclonal antibody, which binds to
10 the C-terminus of PPV-VP1up, reduced PPV PLA₂ activity of VP1up (lane 2), but not that of B19 (lane 4). Adding EGTA to a final concentration of 5 mM abolished PLA₂ activity (lane 5).

B. Sequences outside the conserved pvPLA₂ domain increased pvPLA₂ activity of the PLA₂ motif of expressed PPV-VP1ups. VP1up from M1 to S174 was used throughout this work
15 and was assigned the relative specific activity of 1.0. Sequences within this expressed peptide but outside the PLA₂ motif (dark-grey) contributed to the activity as shown by the relative specific activities.

C. The impact of Ca²⁺ concentration, pH and different substrates on the activity of pvPLA₂. Since 1 mM Ca²⁺, 50 mM Tris-HCl, pH 8.0, and phosphatidylcholine (PC) were used in
20 standard assays, these were set at 100% relative activity. PE = phosphatidylethanolamine, PI = phosphatidylinositol. Samples were measured in triplicate.

D. PLA₂ activity of untreated 0.2 µg virions (lane 1) and after dissociation (lanes 2 & 3) and heat shock (lanes 4 & 5). Bee venom was included as a positive control (lanes 6 & 7). Lane 8 contains the negative control. Samples in lanes 3, 5 and 7 were treated with anti-VP1up
25 antibody.

Figure 4 depicts the predicted 3D-structure of the vPLA₂ domain. The predicted 3D-structure of the vPLA₂ domain, as shown for PPV/B19, showed sequence homology with group III sPLA₂ at the – and C-termini, whereas the centre section resembled group I/II sPLA₂ (sequence of IB pancreatic sPLA₂ shown). The top line shows the position of helices (H) for group I/I, which are inferred from the homologous porcine pancreatic PLA₂. The bottom line shows the position of helices (H) for group III PLA₂. The predicted three-dimensional model of B19/PPV PLA₂ was obtained by homology modeling using the program MOE. Both the three-dimensional structure of bee venom PLA₂ (Group III, in magenta) and the three-dimensional structure of the pancreatic porcine PLA₂ (Group I/II, in orange) were used for homology modeling of the vPLA₂ (centre). Note that residues L, V, I as well as residues Y, F, W were considered equivalent residues. The three-dimensional images were generated with the program GRASP. The color code in the sequence matches that in the vPLA₂ structure. Residues coloured green in vPLA₂ indicate residues that are conserved in both group I/II and group III PLA₂s; orange-coloured residues are conserved in group I/II; and magenta-coloured residues are conserved in group III PLA₂s. The amino acids identified in the vPLA₂ structure were submitted to site-directed mutagenesis.

Figure 5 depicts immunofluorescence of infected cells using wild-type (wt) or mutant virus, with or without NH₄Cl treatment. Immunofluorescence was used to follow the entry and infection of PT cells by the wt or mutant virus. At 0 hours of entry, a strong staining, often patchy, was observed on the cellular membrane. After 4 hours, virus entry led to a cytoplasmic, granular, and perinuclear staining, whereas with NH₄Cl treatment, the virus was still mostly distributed throughout the cytoplasm. After 8 hours, all incoming virus was perinuclear except for wt without NH₄Cl, for which some nuclei became positive. At 12 hours, only wt without NH₄Cl yielded significant numbers of positive nuclei. Some positive nuclei appeared at 18 hours for wt with NH₄Cl, and for the P21W or P21L mutants. The delivery of the viral genome can be seen to be impaired in the case of the mutant virus, and is further inhibited by treatment with NH₄Cl.

Figure 6 shows the binding (A) and entry (B) of ³⁵S-labelled porcine Parvovirus using 4x10⁵ cells and 5x10⁹ wild type (wt) virus particles. Equivalent amounts of mutant viruses were

used for the binding studies, and twice as much for the entry studies (experiments in quadruplicate).

C. ^{35}S -labeled PPV particles (wt or mutant) were added to 2×10^7 cells in Petri dishes and about 20% was recovered in the cells. After cellular fractionation, the distribution of virus
5 over cytosolic and nuclear fractions was measured by liquid scintillation. The relative radioactivity is calculated as follows: [nuclear fraction-bound activity]/[cellular fraction-bound activity]. Standard errors were calculated from three experiments.

D. Co-localization of PPV and LAMP-2 was studied by confocal microscopy. Anti-PPV antibodies were labeled with FITC, giving green fluorescence, and anti-LAMP-2 antibodies
10 with Texas Red, giving red fluorescence. Co-localization of PPV and LAMP-2 resulted in yellow staining.

Figure 7 demonstrates distribution of viral DNA in infected cells. Incoming viral DNA was detected in the cytoplasm by *in situ* hybridization 4 (A) and 8 (B) hours post-infection (p.i.), whereas replicating DNA in the nucleus was found at 12 hours p.i. (C). At 12 hours p.i., no
15 DNA could be detected in the nucleus in the case of the HD (D) and P21^{PW} mutants. Aphidicoline, a DNA polymerase inhibitor, also prevented the appearance of viral DNA in the nucleus (F). The sPLA₂ inhibitors tetracain (TC), and oleyloxyethyl phosphorylcholine (OP), also reduced the number of nuclei containing viral DNA (G).

Figure 8 shows nucleic acid sequences of cloned V1up regions of porcine parvovirus [SEQ
20 ID NO: 50], B19 human parvovirus [SEQ ID NO: 52], and *Galleria mellonella* densovirus [SEQ ID NO: 54].

Figure 9 shows amino acid sequences of cloned V1up regions of porcine parvovirus [SEQ ID NO: 49], B19 human parvovirus [SEQ ID NO: 50], and *Galleria mellonella* densovirus [SEQ ID NO: 52].

25 **Figure 10** shows the 5234 base pair nucleic acid sequence of *Acheta domesticus* densovirus [SEQ ID NO: 65].

Figure 11 shows the nucleic acid sequence of the genomic region containing the PLA₂ motif for *Acheta domesticus* densovirus [SEQ ID NO: 64] and *Mythimna loreyi* densovirus [SEQ ID NO: 62].

Figure 12 shows the amino acid sequence of the region containing the PLA₂ motif for *Acheta domesticus* densovirus [SEQ ID NO: 63] and *Mythimna loreyi* densovirus [SEQ ID NO: 61]. The conserved histidine (H) and aspartic acid (D) residues are highlighted.

Figure 13 shows the sequences identified by phage display that bind to viral PLA₂.

DETAILED DESCRIPTION OF THE INVENTION

Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention pertains. For the purposes of the present invention, the following terms and abbreviations are defined below.

The term "agent" is used herein to denote a chemical compound, a mixture of chemical compounds, a biological macromolecule, or an extract made from biological materials such as bacteria, plants, fungi, or animal (particularly mammalian) cells or tissues.

The term "amino acid motif" is used herein to denote a characteristic sequence of amino acids the presence of which can serve to identify members of a family of proteins. It is understood by those of skill in the art that variations of one or two amino acids within a motif does not preclude a protein from belonging to the family defined by the motif.

This invention is based on the unexpected finding that PLA₂ activity is present in the capsids of viruses. This invention is also based on the unexpected determinations that this viral PLA₂ activity is essential for infection and that inhibition of this PLA₂ activity decreases the infectivity and/or replication of the viruses.

Characterization of Viral PLA₂s

The present invention describes a class of PLA₂s found in viruses. These viral PLA₂s have significant differences from known PLA₂s, including structure (Figure 4), biological function, and amino acid sequences (Figure 2); thus, these viral PLA₂s form a new group of PLA₂s
5 within the PLA₂ superfamily.

The present invention provides viral PLA₂s containing a unique, characteristic sequence profile. Viral PLA₂s comprise the amino acid motif [WY]-[CIVL]-G-x-[GP]-[GKNS]-x(8,14)-D-x(2)-[AC]-x(2)-H-D-x(2)-Y, where the notation [WY] represents alternative amino acids at this position, x is any amino acid, x(2) represents a stretch of 2 amino acids
10 and x(8,14) represents a stretch of between 8 and 14 amino acids. The above motif is characteristic of viral PLA₂ sequences.

In accordance with one aspect of the present invention, there is provided polynucleotides that encode viral PLA₂s containing the amino acid motif [WY]-[CIVL]-G-x-[GP]-[GKNS]-x(8,14)-D-x(2)-[AC]-x(2)-H-D-x(2)-Y. The present invention further relates to polypeptides
15 which contain the viral PLA₂ amino acid motif and exhibit phospholipase A₂ activity.

In one embodiment of the present invention, the polypeptide is a variant of a naturally occurring viral PLA₂ which may have a substitution, deletion, or addition of one or more amino acids. In a related embodiment, there is provided polynucleotides that encode such variants.

20 The present invention further provides variants of the viral PLA₂s that may be inactive due to the substitution of an active site residue, or that may contain slight variations of the above amino acid motif. Analogous variants exist in the mammalian PLA₂s in the form of pseudogenes. The viral PLA₂s of the present invention may be obtained from any virus presently identified and from viruses yet to be identified or may be prepared by recombinant
25 molecular biology techniques or chemical synthesis.

1) *PLA₂ activity*

The viral PLA₂s of the present invention are ultimately defined by their PLA₂ activity. PLA₂ activity is defined as the ability to hydrolyse the fatty acid from the *sn*-2 position of phospholipids to yield lysophosphatidyl compounds.

- 5 One exemplary assay to determine PLA₂ enzyme activity is described. PLA₂ enzyme activity can be established using the *E. coli* radioactive assay (Elsbach and Weiss (1991) *Methods Enzymol* 197:24). *E. coli* phospholipids are metabolically labeled for 3 hours with [¹⁴C]-oleic acid (95% in *sn*-2 position). The bacteria are then autoclaved for 15min. to inactivate bacterial phospholipases, and washed. The viral sample (1-1000 pg) is added to the bacteria
- 10 (10,000 cpm) in standard buffer (50 mM Tris-HCl, pH 8.0, and 10 mM CaCl₂) and incubated for 30min. at 37°C. The reaction is stopped by the addition of ice-cold 1% BSA. After centrifugation, 50% of the supernatant is counted. PLA₂ activity is identified and/or quantified by the release of radioactive fatty acids from the phospholipids into the supernatant.
- 15 A mixed micelles assay can also be used to establish PLA₂ activity. The substrates used are 6 μM of either L-3-phosphatidylcholine, 1,2-di[1-¹⁴C]oleoyl (specific activity 110 mCi/mmol), L-3-phosphatidylinositol L-1-stearoyl-2-[1-¹⁴C]arachidonoyl (specific activity 48 mCi/mmol), or phosphatidylethanolamine L-1-palmitoyl, 2-[1-¹⁴C]arachidonoyl (specific activity 54.6 mCi/mmol). The assay is based on that described by Manjunath *et al.*, (1994)
- 20 *Biochem J.* 303:121, with the following modifications: 1 mM Triton X-100 is used instead of deoxycholate; the total reaction volume is 50 μl and the reaction is stopped after 10 min by adding 80 μl chloroform/methanol (2:1) and 50 μl of saturated KCl solution. Separation is on silica gel thin layer chromatography (TLC) plates. Optimal separation of the phosphatidylcholine reaction products is obtained using a solvent solution containing
- 25 chloroform, methanol and water at a ratio of 65:35:4. For phosphatidyl ethanolamine and phosphatidylinositol, chloroform and methanol (87:13) are used as a solvent. The separated products are quantified using a Molecular Dynamics PhosphorImager SI after drying. PLA₂ activity is identified and/or quantified by the presence of radioactive fatty acids, released from phospholipids, on each TLC plate.

2) *Parvoviral PLA₂s*

In one embodiment of the present invention, the viral PLA₂ is a parvovirus PLA₂. In a related embodiment the parvovirus PLA₂ is present in the VP1-unique part (VP1up) of the largest capsid protein of parvoviruses.

- 5 In accordance with the present invention, the parvovirus PLA₂s comprise the following, more specific version of the above amino acid motif: Y-x-G-P-G-x(12)-D-x(2)-A-x(2)-H-D-x(2)-Y, where x is any amino acid, x(2) represents a stretch of 2 amino acids, x(12) represents a stretch of 12 amino acids and wherein one of the specified amino acids Y, G, P, G, D, A, D, or Y may be exchanged singly for any other amino acid. This invention includes viral PLA₂s
- 10 obtained from known parvoviruses, including, but not limited to, mammalian parvoviruses, such as canine parvovirus (CPV), mink enteritis virus (MEV), porcine parvovirus (PPV), bovine parvovirus (BPV), goose parvovirus (GPV), feline panleukopenia virus (FPLV), the human parvoviruses, such as B19, V9, and RA-1, and insect parvoviruses, such as *Gm* DNV; *Mi* DNV; *Jc* DNV; *Pi* DNV; *Ds* DNV; *Cp* DNV; *Ce* DNV; *Ad* DNV; *Pf* DNV; and *Bm*
- 15 DNV. The parvoviral PLA₂s of the present invention may also be obtained from parvoviruses yet to be identified. The protein sequences determined for 30 different parvoviral PLA₂s are shown in Figure 2.

Preparation of Viral PLA₂s

- Unless otherwise specified, the viral PLA₂ proteins, peptides or fragments thereof, of the
- 20 present invention are prepared in such a manner that their intrinsic enzymatic activity is retained. Further, amino acid residues may be deleted, added or substituted for those that appear in the amino acid sequences of the viruses of interest in instant invention. It should also be appreciated that the present invention contemplates amino acid sequences that are equivalent to, or constitute active fragments of, the amino acid sequences for the viral PLA₂
- 25 enzymes of the present invention.

The proteins of the present invention can be prepared through the use of recombinant techniques, or from viral extracts. In general, viral PLA₂ proteins according to the instant invention can be produced by transformation (transfection, transduction, or infection) of a host cell with all or part of a viral PLA₂-encoding DNA fragment in a suitable expression
5 vehicle. Suitable expression vehicles include: plasmids, phagemids, viral particles, and phage. The entire expression vehicle, or a part thereof, can be integrated into the host cell genome. In some circumstances, it is desirable to employ an inducible expression vector.

Those skilled in the field of molecular biology will understand that any of a wide variety of expression systems can be used to provide a recombinant viral PLA₂ protein of the instant
10 invention. The precise host cell used is not critical to the instant invention. A viral PLA₂ protein can be produced in a prokaryotic host (e.g., *E. coli* or *B. subtilis*) or in a eukaryotic host (e.g., *Saccharomyces* or *Pichia*; mammalian cells, e.g., COS, NIH 3T3, CHO, BHK, 293, or HeLa cells; or insect cells) using techniques that are standard in the field of molecular biology.

- 15 The methods of transformation or transfection and the choice of expression vehicle will depend on the host system selected. Transformation and transfection methods are described, e.g., in Ausubel *et al.* (1994) Current Protocols in Molecular Biology, John Wiley & Sons, New York. Expression vehicles can be chosen from those provided, e.g. in Cloning Vectors: A Laboratory Manual (Pouwels *et al.*, 1985, Supp. 1987).
- 20 The host cells harbouring the expression vehicle can be cultured in conventional nutrient media adapted as necessary for activation of a chosen gene, repression of a chosen gene, selection of transformants, or amplification of a chosen gene. One example of an expression system is the mouse 3T3 fibroblast host cell transfected with a pMAMneo expression vector (Clontech, Palo Alto, Calif.). pMAMneo provides an RSV-LTR enhancer linked to a
25 dexamethasone-inducible MMTV-LTR promotor, together with an SV40 origin of replication, which allows replication in mammalian systems, a selectable neomycin gene, and SV40 splicing and polyadenylation sites. DNA encoding a viral PLA₂ protein can be inserted into the pMAMneo vector the appropriate orientation to allow expression of the protein. The

recombinant viral PLA₂ protein can then be isolated from the expression system using standard techniques. Other host cells that can be used in conjunction with the pMAMneo expression vehicle include COS cells and CHO cells (ATCC Accession Nos. CRL 1650 and CCL 61, respectively).

- 5 Alternatively, viral PLA₂ polypeptides can be produced as fusion proteins. For example, the expression vector pUR278 (Ruther *et al.* (1983) *EMBO J.* 2, 1791) can be used to create lacZ fusion proteins, or the pGEX vectors can be used to create glutathione S-transferase (GST) fusion proteins. In general, GST-fusion proteins are soluble and can be easily purified from lysed cells by adsorption to glutathione-agarose beads, followed by elution in the presence
10 of free glutathione. The pGEX vectors are designed to provide thrombin or factor Xa protease cleavage sites so that the cloned target gene product can be released from the GST moiety.

- Specific initiation signals may also be required for efficient translation of inserted nucleic acid sequences. These signals include the ATG initiation codon and adjacent sequences. In
15 cases where an entire native viral PLA₂ gene or cDNA, including its own initiation codon and adjacent sequences, is inserted into the appropriate expression vector, no additional translational control signals may be needed. In other cases, exogenous translational control signals, including, perhaps, the ATG initiation codon, must be provided. Furthermore, the initiation codon must be in phase with the reading frame of the desired coding sequence to
20 ensure translation of the entire insert. These exogenous translational control signals and initiation codons can be of a variety of origins, both natural and synthetic. If necessary, the efficiency of expression can be enhanced by the inclusion of appropriate transcription enhancer elements and / or transcription terminators (Bittner *et al.* (1987) *Methods in Enzymol.* 153, 516).

- 25 Alternatively, a viral PLA₂ protein can be produced from a stably-transfected mammalian cell line. A number of vectors suitable for stable transfection of mammalian cells are available to the public, see, *e.g.* Pouwels *et al.* (*supra*). Methods for constructing such cell lines are also publicly available and known to those skilled in the art, see for example Ausubel *et al.*

(*supra*). In one example, cDNA encoding the viral PLA₂ protein can be cloned into an expression vector that includes the dihydrofolate reductase (DHFR) gene. Integration of the plasmid and, therefore, the viral PLA₂ protein-encoding gene into the host cell chromosome can then be selected for by including an appropriate amount of methotrexate in the cell culture medium (as described in Ausubel *et al.*, *supra*). This dominant selection can be accomplished in most cell types.

Fusion proteins can be readily purified by utilising an antibody or other binding partner specific for the fusion protein being expressed. For example, a system described in Janknecht *et al.* (1981) *Proc. Natl. Acad. Sci. USA* 88:8972, allows for the ready purification of non-denatured fusion proteins expressed in human cell lines. In this system, the gene of interest is subcloned into a vaccinia recombination plasmid such that the open reading frame of the gene is translationally fused to an amino-terminal tag consisting of six histidine residues. Extracts from cells infected with the recombinant vaccinia virus are loaded onto a Ni²⁺ nitriloacetic acid-agarose column, and the histidine-tagged proteins are selectively eluted with imidazole-containing buffer.

Alternatively, a viral PLA₂ or a portion thereof, can be fused to an immunoglobulin Fc domain. Such a fusion protein can be readily purified using a protein A column.

It will be apparent to one skilled in the art that the viral PLA₂ peptides disclosed herein can be created by designing nucleic acid sequences that encode for viral PLA₂, but which differ, by reason of the redundancy of the genetic code, from the sequences disclosed herein. Accordingly, the degeneracy of the genetic code further enables major variations in the nucleotide sequence of a nucleic acid molecule but does not broaden the scope of the present invention since the amino acid sequence of the encoded protein remains unchanged. Based upon the degeneracy of the genetic code, variant nucleic acid sequences may be derived from the nucleic acid sequences disclosed herein. These variant nucleic acid sequences can be produced by modifying or synthesizing nucleic acid sequences. Variant nucleic acid sequences include deletion, addition, substitution, or a combination thereof, of different nucleotides.

A viral PLA₂ protein sequence of the present invention can be identified using the criteria set out above and can be chemically synthesized by methods known to those of skill in the art. Such methods include, but are not limited to, exclusive solid phase synthesis, partial solid phase synthesis, fragment condensation or classical solution synthesis (Merrifield (1963) *J. Am. Chem. Soc.* 85:2149; Merrifield (1986) *Science* 232:341). Further, viral PLA₂ peptides can be isolated and purified by standard purification methods including chromatography (e.g. ion exchange, affinity, and sizing column chromatography or high performance liquid chromatography), centrifugation, differential solubility, or by other standard techniques familiar to a worker skilled in the art. Once synthesized, the authenticity of the resulting peptides can be verified using the assay procedures disclosed herein.

Inactive Viral PLA₂s

The present invention also includes viral PLA₂ proteins, peptides or fragments thereof, the sequence of which produces an inactive protein (*i.e.* one that no longer possesses PLA₂ activity), or has been manipulated to produce an inactive protein.

- 15 Once a viral PLA₂ protein sequence has been identified using the criteria set out above, the nucleic acid sequence encoding the protein can be isolated. Standard molecular biology techniques can then be employed to alter the nucleic acid sequence and, thus the protein sequence that it encodes. Methods of altering nucleic acid sequences are well-known in the art, for example, site-directed mutagenesis or PCR-based techniques can be employed.
- 20 Naturally inactive mammalian sPLA₂ proteins, generated from pseudogenes, are known. An amino acid motif: HDXXY (where X = any amino acid) is found in the active site of all known sPLA₂ proteins (Dennis (1994) *J. Biol. Chem.* 269:13057-13060). In the inactive sPLA₂ proteins, the conserved aspartic acid residue (D), subsequent to the conserved histidine residue (H) in the above motif, is frequently mutated to a basic amino acid. The HDXXY motif is also found in viral PLA₂s, therefore, one example of an inactive viral PLA₂ protein
- 25

would be one in which the aspartic acid residue in this motif is exchanged for a basic amino acid such as, lysine, arginine or histidine..

It is envisioned that such inactive viral PLA₂ proteins or peptides will retain their ability to induce a specific immune response and that they will, therefore, be useful in the development
5 of vaccines.

Uses of Viral PLA₂s

The viral PLA₂s of the present invention can be used in manufacturing processes, in the clinical diagnosis and treatment of diseases, and in basic research settings.

1) Research

10 At present, purified bee venom PLA₂ is commonly used in research, for example as a reference standard when determining the PLA₂ enzyme activity contained in various preparations of isolated enzyme. The viral PLA₂ of PPV, however, has a 40 to 100-fold higher specific activity than that of bee venom PLA₂ which makes it better suited to various research applications. In addition, while some parvovirus PLA₂s demonstrate specificity for
15 certain phospholipids (e.g. GmDNV PLA₂ is specific for phosphatidylinositol), others have a broad substrate specificity thus making their use widely applicable.

2) Screening for viral PLA₂ inhibitors

The viral PLA₂s of the present invention are also useful for screening compounds in order to identify agents capable of inhibiting PLA₂ activity. Useful inhibitory agents are identified
20 with a range of assays employing viral PLA₂s or nucleic acids encoding viral PLA₂s. As examples, protein binding assays, nucleic acid binding assays, gel shift assays, cell-based assays, mixed micelle assays, and the like are useful approaches.

In one embodiment of the present invention, there is provided a screening assay in which a candidate inhibitor is added to a solution containing a viral PLA₂ and a labelled phospholipid

substrate, and the inhibition of PLA₂ activity is then determined. In order to determine whether the inhibitory activity of the candidate inhibitor compound is specific, PLA₂ activity in the presence of the candidate inhibitor, is compared to the following: a) the PLA₂ activity in the absence of the candidate compound, b) in the presence of a well-characterized PLA₂ inhibitor, c) in the presence of a PLA₂ agonist, d) in the presence of activators and inhibitors of other phospholipase enzymes, for example PLC, and e) in the presence of both the candidate inhibitor agent and the well-characterized PLA₂ inhibitor. PLA₂ activity can be determined according to the methods described herein. Such *in vitro* screening procedures are especially useful in identifying candidate agents that can be subsequently tested *in vivo* for their ability to inhibit viral PLA₂s.

In another embodiment of the present invention, the *in vitro* assay is an automated, cost-effective, high-throughput screening protocol that can be used to survey a large number of test compounds for their ability to inhibit viral PLA₂ activity.

Potential inhibitory compounds are screened from large libraries of synthetic or natural compounds. Numerous means are currently used for random and directed synthesis of saccharide, peptide, and nucleic acid based compounds. Synthetic compound libraries are commercially available from a number of companies including Maybridge Chemical Co. (Trevillet, Cornwall, UK), Comgenex (Princeton, N.J.), Brandon Associates (Merrimack, N.H.), and Microsource (New Milford, Conn.). A rare chemical library is available from Aldrich (Milwaukee, Wis.). Combinatorial libraries are also available and can be prepared according to standard procedures. Alternatively, libraries of natural compounds in the form of bacterial, fungal, plant, and animal extracts are available from, e.g., Pan Laboratories (Bothell, Wash.) or MycoSearch (North Carolina), or are readily producible. Additionally, natural and synthetically produced libraries and compounds are readily modified through conventional chemical, physical, and biochemical means. The agent library can be naive or can be composed of structural analogs of known PLA₂ inhibitors, or can be a combination of both.

3) Gene Therapy

Viral phospholipase A₂ has been shown to be critical for the transfer of the viral genome from the late endosome to the nucleus (see Figure 7). Therefore, the present invention provides a method of improving viral-based vectors for gene therapy. An example would be the adeno-associated virus (AAV) vector currently envisioned for gene therapy. Inclusion of the viral PLA₂ gene into known gene therapy vectors can optimize the transfer of the therapeutic gene-carrying genome to the nucleus of target cells. If necessary, the viral PLA₂ gene may be modified for this purpose, for example, by site-directed mutagenesis, for optimal effectiveness and specificity.

4) Viral PLA₂-Binding Peptides

The present invention also provides for polypeptides and peptides that bind the viral PLA₂s. One exemplary method of identifying such peptide is by phage display techniques. Phage display libraries of random short peptides are commercially available, e.g. from New England Biolabs, Inc., which are utilized through an *in vitro* selection process known as "panning". In its simplest form, panning involves incubating the library of phage displayed peptides with a plate, or bead, coated with the target molecule, washing away unbound phage particles and finally eluting the specifically bound phage. For the purposes of the present invention, the target molecule is a viral PLA₂, or fragment thereof.

In one embodiment of the present invention, the target molecule is porcine parvovirus PLA₂ and the peptides displayed by phage display library are 12-mers.

The peptide(s) displayed by the specifically-binding phage are then isolated and sequenced by standard techniques known to those of skill in the art. In some instances the binding strength of the isolated peptide is then tested using standard techniques.

In one embodiment of the present invention, there is provided polynucleotides which encode the viral PLA₂-binding peptide. These polynucleotides can be cloned and fused with a heterologous nucleic acid. In a related embodiment, the resulting fusion gene is used to produce a fusion protein comprising the viral PLA₂-binding peptide and a heterologous protein. As discussed herein, a worker skilled in the art would readily appreciate that the

polynucleotides of the present invention can be cloned with the appropriate regulatory sequences for expression.

In one embodiment of the present invention, there is provided a fusion protein comprising a viral PLA₂-binding peptide and thioredoxin. The thioredoxin fusion protein can be used in
5 an ELISA assay for detection of viral PLA₂s. In this assay the viral PLA₂-binding peptide of the fusion protein binds the viral PLA₂ in a sample, and the resulting complexes are detected using anti-thioredoxin antibodies. The antibodies are either labelled directly, or are detected with a secondary antibody that is labelled, for example with an enzyme in the case of ELISA assays. Such an assay is useful for detecting virus in a sample from a subject suspected of
10 having a viral infection.

Furthermore, peptides that bind to the viral PLA₂s can be directly labelled by techniques known in the art and used as probe molecules. As such, labelled viral PLA₂-binding peptides are deemed to be within the scope of those skilled in the art in view of the teaching provided herein.

15 Alternatively, peptides can be selected that, by binding to the viral PLA₂, specifically inhibit the activity of the protein. The inhibitory activity of the peptides is determined using assays as known in the art and as described herein. Inhibitory viral PLA₂-binding peptides can be used therapeutically to reduce or eliminate viral infectivity. One embodiment of the present ideation provides pharmaceutical compositions comprising one or more viral PLA₂-binding
20 peptides or polynucleotides encoding such peptides and a pharmaceutically acceptable diluent or excipient.

Anti-viral Agents

The present invention further provides a class of anti-viral agent that selectively inhibits viral PLA₂ activity. Viral PLA₂ activity is required in order for the virus to enter a host nucleus;
25 thus, inhibition of viral PLA₂ activity blocks the ability of a virus to infect a host cell. Since

the sequence of the viral enzyme is very different from that of the host enzymes, specific inhibitors can be developed that do not impair host PLA₂ enzyme activities. The present invention provides anti-viral PLA₂ agents that inhibit viral PLA₂ activity but that are nontoxic to the host.

5 *1) Selection of Inhibitors (Active Agents)*

The anti-viral PLA₂ agents of the present invention are initially selected on the basis of the agent's capacity for inhibiting viral PLA₂ *in vitro*. Inhibition of PLA₂ activity can be determined using assays that monitor phospholipase activity including the screening assay of the present invention. This initially selected agent is then be administered to a cultured
10 host cell population, which is subsequently exposed to a virus. The capacity of the agent to produce a detectable reduction in the infectivity and / or replication of the virus in the treated cell culture, in comparison to an untreated cell culture, is determined according to standard procedures known in the art, for example by measuring plaque forming units. Agents capable of reducing infectivity and / or replication of the virus are thereby identified as active agents.

15 In addition to exhibiting anti-viral activity, the anti-viral PLA₂ agents must also be selective. A selective anti-viral PLA₂ agent produces a preferential inhibition of viral PLA₂ as compared to inhibition of mammalian or host PLA₂s. Typically, the anti-viral PLA₂ agent required to produce inhibition of 50% of viral PLA₂ catalytic activity is at least one order of magnitude lower than the concentration required to produce inhibition of 50% of the catalytic
20 activity of phospholipases other than viral PLA₂, including mammalian (host) PLA₂, PLA₁, PLC and PLD. Two exemplary methods that can be used to determine the specificity of potential anti-viral PLA₂ agents are described briefly below, although it will be appreciated by those skilled in the art that alternative methods can also be employed.

In one method PLA₂ enzyme activity is measured using a phospholipase assay familiar to one
25 skilled in the art. An example of a suitable phospholipase assay for this procedure is described in detail in Example III. Briefly, isolated virus preparations are homogenized and centrifuged in a manner familiar to one skilled in the art of enzyme purification and supernatants are obtained. PLA₂ activity in the supernatant is then measured in the absence

and presence of various potential anti-viral PLA₂ agents. A similar procedure is used to determine the activity of mammalian or host cell PLA₂s, PLA₁s, PLCs and PLDs, in the absence and presence of the potential anti-viral agents. The selection of an anti-viral PLA₂ agent is determined by comparing the kinetic parameters of both the viral PLA₂ and the host cell PLA₂. An example of a potentially useful anti-viral PLA₂ agent for clinical trials is one that inhibits viral PLA₂ activity by at least 50 % with an IC₅₀ at least one order of magnitude lower than that which inhibits the host cell PLA₂.

A second method to determine the activity of potential anti-viral PLA₂ agents is by immunofluorescence assay as described in Example VII and shown in Figure 5. A preparation of isolated virus is labelled with a fluorescent marker and then administered to a host cell culture. Entry and infection of the host cells by the virus is monitored by tracking this fluorescent tag over a period of time in the presence and absence of a potential anti-viral PLA₂ agent. The ability of the virus to infect the cell, and thus the capacity of an anti-viral PLA₂ agent to inhibit infectivity, is a function of the amount of fluorescence tag that is viewed within the cell. The higher the amount of fluorescence the weaker the inhibitory capacity of the anti-viral PLA₂ agent.

2) Antisense Oligonucleotides

The present invention contemplates antisense oligonucleotides designed to inhibit viral PLA₂ expression, wherein the antisense oligonucleotides can be complementary to all or part of the viral PLA₂ sequences, such as antisense oligonucleotides to the viral PLA₂ gene or mRNA. Such complementary antisense polynucleotides may include nucleotide substitutions, additions, deletions, or transpositions, so long as specific hybridization to the relevant target sequence is retained as a functional property of the polynucleotide. Antisense oligonucleotides include soluble antisense RNA or DNA oligonucleotides which can hybridize specifically to viral PLA₂ mRNA species and prevent transcription of the mRNA species and/or translation of the encoded polypeptide (Ching *et al.*, (1989) *Proc. Natl. Acad. Sci. U.S.A.* 86:10006; Broder *et al.*, (1990) *Ann. Int. Med.* 113:604; Loreau *et al.*, (1990) *FEBS Letters* 274:53; Holcenberg *et al.*, WO91/11535; U.S. Ser. No. 07/530,165; WO91/09865; WO91/04753; WO90/13641; and EP 386563). In one embodiment of the

present invention, the antisense oligonucleotides of the present invention exhibit little, or no, inhibition of expression of non-viral PLA₂s.

Antisense molecules are generally targeted to specific nucleic acids. "Targeting" an antisense compound to a particular nucleic acid, in the context of this invention, is a multistep process.

- 5 The process usually begins with the identification of a nucleic acid sequence whose function is to be modulated. In the present invention, the target is a nucleic acid molecule encoding the viral PLA₂ gene. The targeting process also includes determination of a site or sites within this gene for the antisense interaction to occur such that the desired effect, e.g., detection or modulation of expression of the protein, will result. Within the context of the present
- 10 invention, examples of intragenic sites for antisense interaction are the regions encompassing the translation initiation and the termination codon of the open reading frame (ORF) of the gene. The terms "translation initiation codon" or "start codon" can encompass several codon sequences. In the context of the invention, "start codon" and "translation initiation codon" refer to the codon, or codons, that may be present *in vivo* to initiate translation of an mRNA
- 15 molecule transcribed from a gene encoding a viral PLA₂ gene, regardless of the sequence(s) of such codons.

- The open reading frame (ORF) or "coding region," which is known in the art to refer to the region between the translation initiation codon and the translation termination codon, also can be targeted effectively. Other target regions include the 5' untranslated region (5'UTR)
- 20 and the 3' untranslated region (3'UTR). The 5'UTR is known in the art to refer to the portion of an mRNA in the 5' direction from the translation initiation codon, and thus includes nucleotides between the 5' cap site and the translation initiation codon of an mRNA, or corresponding nucleotides on the gene. The 3'UTR is known in the art to refer to the portion of an mRNA in the 3' direction from the translation termination codon, and thus includes
- 25 nucleotides between the translation termination codon and 3' end of an mRNA, or corresponding nucleotides on the gene. The 5' cap of an mRNA comprises an N⁷-methylated guanosine residue joined to the 5'-most residue of the mRNA via a 5'-5' triphosphate linkage. The 5' cap region of an mRNA is considered to include the 5' cap structure itself, as well as

the first 50 nucleotides adjacent to the cap. The 5' cap region, therefore, can also be a target region for the antisense molecule.

- Antisense oligonucleotides are chosen that are sufficiently complementary to the target, i.e., hybridize sufficiently well and with sufficient specificity, to give the desired effect. In the context of this invention, "hybridization" means hydrogen bonding, which may be Watson-Crick, Hoogsteen or reversed Hoogsteen hydrogen bonding, between complementary nucleoside or nucleotide bases. Adenine and thymine, for example, are complementary nucleobases which pair through the formation of hydrogen bonds. "Complementary," as used herein, refers to the capacity for precise pairing between two nucleotides: if a nucleotide at a certain position of an oligonucleotide is capable of hydrogen bonding with a nucleotide at the same position of a DNA or RNA molecule, then the oligonucleotide and the DNA or RNA are considered to be complementary to each other at that position. The oligonucleotide and the DNA or RNA are complementary to each other when a sufficient number of corresponding positions in each molecule are occupied by nucleotides that can hydrogen bond with each other such that stable and specific binding occurs between the oligonucleotide and the DNA or RNA target. It is understood in the art that the sequence of an antisense compound need not be 100% complementary to that of its target nucleic acid to be specifically hybridizable. An antisense compound is specifically hybridizable when binding of the compound to the target DNA or RNA molecule interferes with the normal function of the target DNA or RNA to cause a loss of utility, and when there is a sufficient degree of complementarity to avoid non-specific binding of the antisense compound to non-target sequences under conditions in which specific binding is desired, i.e., under physiological conditions in the case of *in vivo* assays or therapeutic treatment, and in the case of *in vitro* assays, under conditions in which the assays are performed.
- In the context of this invention, the term "oligonucleotide" refers to an oligomer or polymer of ribonucleic acid (RNA) or deoxyribonucleic acid (DNA) or mimetics thereof. This term includes oligonucleotides composed of naturally-occurring nucleobases, sugars and covalent internucleoside (backbone) linkages as well as oligonucleotides having non-naturally-occurring portions which function similarly. Such modified or substituted oligonucleotides

are often preferred over native forms because of desirable properties such as, for example, enhanced cellular uptake, enhanced affinity for nucleic acid target and increased stability in the presence of nucleases.

Examples of modified or substituted antisense compounds useful in this invention include

5 oligonucleotides containing modified backbones or non-natural internucleoside linkages. As defined in this specification, oligonucleotides having modified backbones include those that retain a phosphorus atom in the backbone and those that do not have a phosphorus atom in the backbone. For the purposes of this specification, and as sometimes referenced in the art, modified oligonucleotides that do not have a phosphorus atom in their internucleoside

10 backbone can also be considered to be oligonucleosides. Other oligonucleotide mimetics include those in which both the sugar and the internucleoside linkage of the nucleotide units are replaced with novel groups, while the base units are maintained for hybridization with the target nucleic acid. One such oligonucleotide mimetic that has been shown to have excellent hybridization properties is referred to as a peptide nucleic acid (PNA). In PNA compounds,

15 the sugar-backbone of an oligonucleotide is replaced with an amide-containing backbone, in particular an aminoethylglycine backbone, see for example, Nielsen *et al.*, *Science* (1991) 254:1497.

Modified oligonucleotides containing one or more substituted sugar moieties and / or one or more nucleobase substitutions are also comprehended by the present invention. The present

20 invention also includes antisense compounds which are chimeric compounds. "Chimeric" antisense compounds or "chimeras," in the context of this invention, are antisense compounds, particularly oligonucleotides, which contain two or more chemically distinct regions, each made up of at least one monomer unit, *i.e.* one nucleotide in the case of an oligonucleotide compound. Chimeric oligonucleotides typically contain at least one region

25 wherein the oligonucleotide is modified so as to confer upon the oligonucleotide increased resistance to nuclease degradation, increased cellular uptake, and / or increased binding affinity for the target nucleic acid.

The antisense compounds used in accordance with this invention typically comprise from about 8 to about 100 nucleobases. In one embodiment of the present invention, the antisense compounds comprise from about 8 to about 50 nucleobases. In a related embodiment, the antisense compounds comprise from about 8 to about 30 nucleobases. In another related
5 embodiment, the antisense compounds comprise from about 15 to about 30 nucleobases. The antisense compounds can be conveniently and routinely made through the well-known technique of solid phase synthesis. Equipment for such synthesis is sold by several vendors including, for example, Applied Biosystems (Foster City, CA). Other means for such synthesis known in the art can be additionally or alternatively employed. Similar techniques
10 using phosphorothioates and alkylated derivatives have been employed to produce oligonucleotides.

The antisense compounds used in accordance with the present invention can be synthesized *in vitro*. The present invention also includes genetic vector constructs designed to direct the *in vivo* synthesis of antisense molecules.

15 3) Antibodies

Another example of anti-viral PLA₂ agents are antibodies raised against specific epitopes of viral PLA₂. Viral PLA₂s are significantly different from other PLA₂s in both their sequences and structure. Antibodies, therefore, can be raised against specific viral PLA₂ epitopes to produce PLA₂ inhibitors that are specific to viral PLA₂s and that do not interact with, or
20 inhibit the activity of, non-viral PLA₂.

For the production of antibodies, various hosts including goats, rabbits, rats, mice, humans, and others can be immunized by injection with viral PLA₂ protein, or with any fragment or oligopeptide thereof that has immunogenic properties. Depending on the host species, various adjuvants may be used to increase immunological response. Such adjuvants include, but are
25 not limited to, Freund's, mineral gels such as aluminum hydroxide, and surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, Keyhole limpet hemolysin (KLH), and dinitrophenol. Examples of adjuvants used in humans include, BCG (bacilli Calmette-Guerin) and *Corynebacterium parvum*.

The oligopeptides, peptides, or fragments used to induce antibodies to viral PLA₂ can have an amino acid sequence consisting of as little as about 5 amino acids. In one embodiment of the present invention, amino acid sequences of at least about 10 amino acids are used. These oligopeptides, peptides, or fragments can be identical to a portion of the amino acid sequence
5 of the natural protein that contains the entire amino acid sequence of a small, naturally occurring molecule. If required, short stretches of viral PLA₂ amino acids can be fused with those of another protein, such as KLH, and antibodies to the chimeric molecule can be produced.

Monoclonal antibodies to viral PLA₂ can be prepared using techniques that provide for the
10 production of antibody molecules by continuous cell lines in culture. These include, but are not limited to, the hybridoma technique, the human B-cell hybridoma technique, and the EBV-hybridoma technique (See, e.g., Kohler, G. *et al.* (1975) *Nature* 256:495-497; Kozbor, D. *et al.* (1985) *J. Immunol. Methods* 81:31-42; Cote, R. J. *et al.* (1983) *Proc. Natl. Acad. Sci.* 80:2026-2030; and Cole, S. P. *et al.* (1984) *Mol. Cell Biol.* 62:109-120.)

15 For example, the monoclonal antibodies according to the invention can be obtained by immunizing animals, such as mice or rats, with purified viral PLA₂. Spleen cells isolated from the immunized animals are then immortalized using standard techniques. Those isolated immortalized cells whose culture supernatant contains an antibody that causes an inhibition of the activity of viral PLA₂ with an IC₅₀ of less than 100 ng/ml are then selected and cloned
20 using techniques that are familiar and known to one skilled in the art. The monoclonal antibodies produced by these clones are then isolated according to standard protocols.

The immortalization of the spleen cells of the immunized animals can be carried out by fusing these cells with a myeloma cell line, such as P3X63-Ag 8.653 (ATCC CRL 1580)
25 according to the method in (1980) *J. of Imm. Meth.* 39:285-308. Other methods known to a person skilled in the art can also be used to immortalize spleen cells. In order to detect immortalized cells that produce the desired antibody against the viral PLA₂, a sample of the culture supernatant is tested using an ELISA assay for reactivity with the viral PLA₂. In order to obtain those antibodies that inhibit the enzymatic activity of viral PLA₂, the culture

supernatant of clones that produce antibodies that bind to viral PLA₂ is additionally examined for inhibition of PLA₂ activity using an appropriate assay, such as those described herein. Those clones whose culture supernatant shows the desired inhibition of viral PLA₂ activity are expanded and the antibodies produced by these clones are isolated according to known
5 methods.

In addition, techniques developed for the production of "chimeric antibodies," such as the splicing of mouse antibody genes to human antibody genes to obtain a molecule with appropriate antigen specificity and biological activity, can be used (Morrison, S. L. *et al.* (1984) *Proc. Natl. Acad. Sci.* 81:6851-6855; Neuberger, M. S. *et al.* (1984) *Nature*
10 312:604-608; and Takeda, S. *et al.* (1985) *Nature* 314:452-454). Alternatively, techniques described for the production of single chain antibodies can be adapted, using methods known in the art, to produce viral PLA₂-specific single chain antibodies. Antibodies with related specificity, but of distinct idiotypic composition, can be generated by chain shuffling from random combinatorial immunoglobulin libraries (see for example, Burton D. R. (1991) *Proc.*
15 *Natl. Acad. Sci.* 88:10134-10137).

Antibodies can also be produced by inducing *in vivo* production in the lymphocyte population or by screening immunoglobulin libraries or panels of highly specific binding reagents as disclosed in the literature (Orlandi, R. *et al.* (1989) *Proc. Natl. Acad. Sci.* 86: 3833-3837; Winter, G. *et al.* (1991) *Nature* 349:293-299).

20 Antibody fragments which contain specific binding sites for viral PLA₂ can also be generated. For example, such fragments include, but are not limited to, F(ab')₂ fragments produced by pepsin digestion of the antibody molecule and Fab fragments generated by reducing the disulfide bridges of the F(ab')₂ fragments. Alternatively, Fab expression libraries can be constructed to allow rapid and easy identification of monoclonal Fab fragments with the
25 desired specificity (see for example, Huse, W. D. *et al.* (1989) *Science* 246:1275-1281).

Various immunoassays can be used for screening to identify antibodies having the desired specificity. Numerous protocols for competitive binding or immunoradiometric assays using

either polyclonal or monoclonal antibodies with established specificities are well known in the art. Such immunoassays typically involve the measurement of complex formation between viral PLA₂ and its specific antibody. Examples of such techniques include enzyme-linked immunosorbent assays (ELISAs), radioimmunoassays (RIAs), and
5 fluorescence activated cell sorting (FACS). Alternatively, two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering viral PLA₂ epitopes, or a competitive binding assay can be used (see Maddox, D. E. *et al.* (1983) *J. Exp. Med.* **158**:1211-1216). These and other assays are well known in the art (see for example, Hampton, R. *et al.* (1990) *Serological Methods: A Laboratory Manual*, APS Press, St Paul,
10 Minn., Section IV; Coligan, J. E. *et al.* (1997, and periodic supplements) *Current Protocols in Immunology*, Greene Pub. Associates and Wiley-Interscience, New York, N.Y.; Maddox, D. E. *et al.* (1983) *J. Exp. Med.* **158**:1211-1216).

Use of Anti-Viral PLA₂ Inhibitors

The identified anti-viral PLA₂ agents of the present invention can be used as commercial
15 reagents as standards in toxicological or pharmaceutical evaluations related to PLA₂s.

The identified anti-viral PLA₂ agents of the present invention can also be used in therapeutic applications in animals and humans. The present invention describes methods of decreasing the infectivity and/or replication of a virus in a subject by administering an anti-viral PLA₂ agent to the subject. In one embodiment, the virus is a parvovirus, including mammalian
20 parvoviruses, such as canine parvovirus (CPV), mink enteritis virus (MEV), porcine parvovirus (PPV), bovine parvovirus (BPV), goose parvovirus (GPV), feline panleukopenia virus (FPLV), and human parvoviruses, including adeno-associated virus, B19, V9 (Nguyen *et al.*, (1999) *J. Clin. Microbiol.* **37**:2483), and RA-1 (Stierle *et al.*, (1987) *Ann. Rheum. Dis.* **46**:219); and insect parvoviruses including densovirus for the following hosts: *Gm*
25 *Galleria mellonella*; *Ml Mythimna loreyi*; *Jc Junonia coenia*; *Pi Pseudoplusia includens*; *Ds Diatraea saccharalis*; *Cp Culex pipiens*; *Ce Casphalia extranea*; *Ad Acheta domesticus*; *Pf Periplaneta fuliginos*; and *Bm Bombyx mori*.

Anti-viral PLA₂ agents that selectively block viral PLA₂ activity, including chemical compounds, antisense oligonucleotides, and antibodies, can be used in the methods of the present invention.

1) Antisense Oligonucleotides

- 5 The antisense oligonucleotides of the present invention inhibit production of the viral PLA₂ polypeptides, thus preventing the virus from infecting the host. Compositions containing a therapeutically effective dosage of viral PLA₂ antisense oligonucleotides can be administered for treatment of viral infections.

- 10 Viral PLA₂ antisense compounds can be used as research reagents and diagnostics. As an example, antisense oligonucleotides, which are able to inhibit gene expression with exquisite specificity, may be used to elucidate the function of particular genes in the viral cycle. Antisense compounds can also be used to distinguish between functions of various members of viral biological pathways.

- 15 The specificity and sensitivity of antisense is also harnessed by those of skill in the art for therapeutic uses. Antisense oligonucleotides have been employed as therapeutic moieties in the treatment of disease states in animals and man. Antisense oligonucleotides have been safely and effectively administered to humans and numerous clinical trials are presently underway. It is thus established that oligonucleotides can be useful therapeutic modalities that can be configured to be useful in treatment regimes for treatment *in vitro* (*i.e.* in cell and
20 tissue culture) and *in vivo* (*i.e.* in whole animals, especially humans).

- The present invention also includes pharmaceutical compositions and formulations which include the antisense compounds of the present invention. In one embodiment of the present invention, there is provided a composition comprising one or more antisense oligonucleotides and a diluent. In an alternative embodiment the composition additionally comprises one or
25 more compounds designed to target the antisense oligonucleotide to the affected tissue.

The pharmaceutical compositions of the present invention may be administered in a number of ways depending upon whether local or systemic treatment is desired and upon the area to be treated. Administration may be topical (including ophthalmic and to mucous membranes including vaginal and rectal delivery); pulmonary, *e.g.* by inhalation or insufflation of
5 powders or aerosols, including by nebulizer; intratracheal; intranasal; epidermal and transdermal; oral or parenteral. Parenteral administration includes intravenous, intraarterial, subcutaneous, intraperitoneal or intramuscular injection or infusion; or intracranial, *e.g.*, intrathecal or intraventricular, administration.

2) Administration of Purified or Synthetic Anti-Viral PLA₂ Inhibitors and Antibodies

10 The anti-viral PLA₂ agents of the present invention can be delivered alone or in combination, and can be delivered along with a pharmaceutically acceptable vehicle. Ideally, such a vehicle would enhance the stability and/or delivery properties. The invention also provides for pharmaceutical compositions containing the active factor or fragment or derivative thereof, which can be administered using a suitable vehicle such as liposomes, microparticles
15 or microcapsules. In various embodiments of the invention, it may be useful to use such compositions to achieve sustained release of the active component.

When the anti-viral PLA₂ agents of the present invention are employed for the treatment of viral infections in a subject, they can be formulated into oral dosage forms such as tablets, capsules and the like. The compounds can be administered alone or in combination with
20 conventional carriers, such as magnesium carbonate, magnesium stearate, talc, sugar, lactose, pectin, dextrin, starch, gelatin, tragacanth, methylcellulose, sodium carboxymethyl-cellulose, low melting wax, cocoa butter and the like. Diluents, flavoring agents, solubilizers, lubricants, suspending agents, binders, tablet-disintegrating agents and the like can also be employed. The compounds can be encapsulated with or without other carriers. In all cases,
25 the proportion of active ingredients in said compositions both solid and liquid will be at least to impart the desired activity thereto on oral administration. The anti-viral PLA₂ agents can also be injected parenterally, in which case they are used in the form of a sterile solution containing other solutes, for example, enough saline or glucose to make the solution isotonic.

For administration by inhalation or insufflation, the anti-viral PLA₂ agents can be formulated into an aqueous or partially aqueous solution, which can then be utilized in the form of an aerosol.

For topical use, the anti-viral PLA₂ agents can be formulated in the form of dusting powders,
5 creams or lotions in pharmaceutically acceptable vehicles, which are applied to the affected portion of the skin.

The dosage requirements of the anti-viral PLA₂ agents will vary with the particular compositions employed, the route of administration, the severity of the symptoms presented and the particular subject being treated. Such dosage requirements can be determined by
10 standard clinical techniques, known to a worker skilled in the art. Treatment can be initiated with small dosages less than the optimum dose of the compound and thereafter increased until the optimum effect under the circumstances is reached. In general, the anti-viral PLA₂ agents of the present invention are most administered at a concentration that will afford effective results without causing any harmful or deleterious side effects. The anti-viral PLA₂
15 agents can be either administered as a single unit dose, or the dosage can be divided into convenient subunits administered at suitable times throughout the day.

3) Vaccines

Compositions for use as a vaccine against a virus generally comprise an immunizing amount of a viral PLA₂, or an immunogenic fragment thereof, as an antigen in a pharmaceutically
20 acceptable vehicle. The present invention also provides for inactive viral PLA₂ proteins, peptides, or fragments thereof, which retain their ability to induce a specific immune response, for use as vaccines. For example, such vaccines can be prepared as injectables, such as liquid solutions or suspensions. Solid forms for solubilization or resuspension in a liquid before injection also can be prepared, or the preparation can be emulsified. The active or
25 inactive antigenic capsids for vaccination can be mixed with excipients that are pharmaceutically acceptable and compatible with the active capsids. Examples of suitable excipients are water, saline, dextrose, glycerol, ethanol, or the like, and combinations thereof. In addition, if desired, the vaccine can contain minor amounts of auxiliary substances such

as wetting or emulsifying agents, pH buffering agents, or adjuvants, such as aluminum hydroxide, which enhance the effectiveness of the vaccine. The vaccines are conventionally administered parenterally by injection, for example, subcutaneously or intramuscularly.

The present invention further provides for inactive recombinant live virus vaccines. Methods
5 of generating recombinant virus particles are well-known in the art. Since viral phospholipase A₂ has been shown to be critical for the transfer of the viral genome from the late endosome to the nucleus, recombinant virus particles containing a genetically engineered PLA₂ gene encoding an inactive PLA₂ protein, will be non-infective. Such recombinant virus particles could, therefore, be combined with a pharmaceutically acceptable vehicle as
10 described above and used to immunize a subject against subsequent infection with the active form of the virus.

The vaccines are administered in a manner compatible with the dosage formulation, and in such amount as will be therapeutically effective and immunogenic. The quantity to be administered depends on the subject to be treated, the capacity of the subject's immune
15 system to synthesize antibodies, and the degree of protection desired. Precise amounts of active ingredient required to be administered depend on the judgment of the practitioner and are peculiar to the subject. Suitable regimes for initial administration and booster shots are also variable, but are typified by an initial administration followed in one or two week intervals by a subsequent injection or other administration.

20 4) Gene Therapy

A viral PLA₂ peptide or biologically active fragments thereof, may also be employed in accordance with the present invention by expression of such proteins *in vivo*, which is often referred to as "gene therapy."

Thus, for example, cells from a patient may be engineered with a polynucleotide (DNA or
25 RNA) encoding a polypeptide *ex vivo*, with the engineered cells then being provided to a patient to be treated with the polypeptide. Such methods are well-known in the art. For

example, cells may be engineered by procedures known in the art by use of a retroviral particle containing RNA encoding viral PLA₂ or a biologically active fragment thereof.

Similarly, cells may be engineered *in vivo* for expression of a polypeptide *in vivo* by, for example, procedures known in the art. As known in the art, a producer cell for producing a
5 retroviral particle containing RNA encoding viral PLA₂, or a biologically active fragment thereof, may be administered to a patient for engineering cells *in vivo* and expression of the polypeptide *in vivo*. These and other methods for administering viral PLA₂, or a biologically active fragment thereof, by such method should be apparent to those skilled in the art from the teachings of the present invention. For example, the expression vehicle for engineering
10 cells may be other than a retrovirus, for example, an adenovirus which may be used to engineer cells *in vivo* after combination with a suitable delivery vehicle.

Within the expression vehicle, the nucleic acid sequence encoding the polypeptide of the present invention is under the control of one or more suitable promoters. Suitable promoters which can be employed include, but are not limited to, adenoviral promoters, such as the
15 adenoviral major late promoter; or heterologous promoters, such as the cytomegalovirus (CMV) promoter; the respiratory syncytial virus (RSV) promoter; inducible promoters, such as the MMT promoter, the metallothionein promoter; heat shock promoters; the albumin promoter; the ApoAI promoter; human globin promoters; viral thymidine kinase promoters, such as the Herpes Simplex thymidine kinase promoter; retroviral LTRs (including the
20 modified retroviral LTRs hereinabove described); the SV40 promoter; the ²-actin promoter; and human growth hormone promoters. Alternatively, the promoter can be the native PLA₂ gene promoter.

When the expression vehicle is a retrovirus, a retroviral plasmid vector is employed to transduce packaging cell lines to form producer cell lines. Examples of packaging cells which
25 may be transfected include, but are not limited to, the PE501, PA317, È-2, È-AM, PA12, T19-14X, VT-19-17-H2, ÈCRE, ÈCRIP, GP+E-86, GP+envAm12, and DAN cell lines as described in Miller, *Human Gene Therapy*, Vol. 1, pgs. 5-14 (1990). The vector can transduce the packaging cells through any means known in the art. Such means include, but

are not limited to, electroporation, the use of liposomes, and CaPO_4 precipitation. In one alternative, the retroviral plasmid vector can be encapsulated into a liposome, or PTH to a lipid, and then administered to a host.

The transduced producer cell line generates infectious retroviral vector particles that include
5 the nucleic acid sequence(s) encoding the polypeptides. Such retroviral vector particles then can be employed to transduce eukaryotic cells, either *in vitro* or *in vivo*. The transduced eukaryotic cells will express the nucleic acid sequence(s) encoding the polypeptide. Eukaryotic cells which can be transduced include, but are not limited to, embryonic stem cells, embryonic carcinoma cells, hematopoietic stem cells, hepatocytes, fibroblasts,
10 myoblasts, keratinocytes, endothelial cells, and bronchial epithelial cells.

The viral vector employed can, in one embodiment, be an adenoviral vector that includes essentially the complete adenoviral genome (Shenk *et al.*, (1984) *Curr. Topics Microbiol. Immun.* 111(3):1-39). Alternatively, the viral vector can be a modified adenoviral vector in which at least a portion of the adenoviral genome has been deleted. The viruses used in the
15 construction of viral vectors are generally rendered replication-defective to remove the undesirable effects of viral replication on the target cells.

The DNA sequences encoding the anti-viral PLA_2 agents employed in the present invention can be either cDNA or genomic DNA. DNA encoding either the entire anti-viral PLA_2 agent, or a portion thereof, can be used. Due to the degeneracy of the genetic code, other DNA
20 sequences that encode substantially the same anti-viral PLA_2 agent or a functional equivalent can also be used. Multiple gene copies can also be used.

In order to produce the gene constructs of the invention, recombinant DNA and cloning methods, which are well known to those skilled in the art, can be utilized (see Sambrook *et al.*, *Molecular Cloning, A Laboratory Manual*, 2d ed. (New York: Cold Spring Harbor
25 Laboratory Press, 1989), including the use of restriction enzymes, site-directed mutagenesis, ligation, homologous recombination, and transfection techniques.

It will be appreciated that administration of the viral vectors of the present invention for gene therapy will be by procedures well-established in the pharmaceutical arts, *e.g.* by direct delivery to the target organ, tissue or site, intranasally, intravenously, intramuscularly, subcutaneously, intradermally and through oral administration, either alone or in
5 combination.

The dosages administered will vary from subject to subject and will be determined by the level of decrease of infectivity and/or replication of the particular virus balanced against any risk or deleterious side effects.

- 10 The methods of the present invention will be particularly useful when the subject is infected by a particular virus. As an example, parvovirus B19 infection may lead to fifth disease in normal individuals, transient aplastic crisis in patients with underlying hemolysis, and chronic anemia due to persistent infection in immunocompromised patients. B19 infection in pregnancy can lead to hydrops fetalis and fetal loss and/or congenital infection. B19 has
15 also been associated with inflammation and autoimmune diseases, such as rheumatoid arthritis (RA), systemic lupus erythematosus (SLE), adult-onset Still's disease, and polyarthritis. It is envisioned, therefore, that the anti-viral PLA₂ agents of the present invention can be used to decrease the infectivity and/or replication of B19 in subjects with these disorders.

20 *Improvement over Current Tools*

- The viral PLA₂s of the present invention are an improvement over PLA₂ standards used currently in manufacturing, clinical, and research settings. At present, bee venom is commonly used as a standard in measuring PLA₂ activity. The viral PLA₂ of PPV, however, has a 41-fold higher specific activity than bee venom. This high activity makes PPV PLA₂
25 ideal for use as a PLA₂ standard. The present invention overcomes the shortcomings and drawbacks often associated with treatment of viral infections, through the identification of a novel family of viral PLA₂ genes and nucleic acid sequences, amino acid sequences, clones,

vectors, antisense nucleotide sequences, and cell lines. The improvement over current tools lies in the aspects of the present invention that can be used to specifically inhibit the infectivity and / or replication of many pathogenic viruses known to cause disease in humans. The instant application describes for the first time the inhibition of a newly characterized
5 class of phospholipase A₂ enzyme found in many viruses. Inhibition of these viral-specific enzymes is less toxic to the host since host-specific enzymes are not inhibited. This specificity is largely due to the significant differences in the sequences of viral versus non-viral PLA₂s. Thus, the present invention provides a powerful new anti-viral tool, the use of which will not affect host PLA₂s and phospholipase enzymes, but will dramatically decreases
10 the viral PLA₂. These viral specific PLA₂s are critical components for the replication of the virus and may further be required for the pathologies associated with viral infection.

The present invention is described in further detail in the following non-limiting examples. It is to be understood that the examples described below are not meant to limit the scope of the present invention. It is expected that numerous variants will be obvious to the person
15 skilled in the art to which the present invention pertains, without any departure from the spirit of the present invention.

EXAMPLES

Materials and Methods

Expression of VP1up fragments:

20 Expressed VP1ups were obtained as follows: the pBADTBX vector was constructed from the pBAD/TOPO Thiofusion expression vector (Invitrogen) by inserting a TEV protease site followed by a polylinker (*Bgl*II, *Xba*I) downstream of the thioredoxin gene. The vector contains the sequence for a His-tag downstream of this sequence. VP1up regions from porcine parvovirus [SEQ ID NO: 50], B19 human parvovirus [SEQ ID NO: 52], *Galleria*
25 *mellonella* densovirus [SEQ ID NO: 54], *Bombyx mori* densovirus [SEQ ID NO: 58], *Casphalia extranea* densovirus [SEQ ID NO: 56] and adeno-associated virus [SEQ ID NO: 60] were cloned into the *Bgl*II-*Xba*I sites to obtain the respective fusion proteins. These

proteins could be purified using the His-tag on a Ni-agarose column. Subsequent cleavage by TEV protease removed the thioredoxin moiety from the fusion protein. PLA₂ activity of the fusion protein, however, was not significantly affected by the presence of thioredoxin. Usual yields were about 1 mg/litre.

5 Polyclonal Anti-V1up Antibodies:

Using the above expression vectors containing the V1up regions, PLA₂ was expressed and purified as described. About 250 mg of the fusion protein was injected into rabbits with adjuvants (subcutaneously and intramuscularly) according to standard protocols. Immunization was repeated after three weeks. The polyclonal antibodies that were generated
10 were isolated by standard techniques.

Assaying PLA₂ Activity:

PLA₂ enzyme activity was established using the *E. coli* radioactive assay (Elsbach and Weiss (1991) *Methods Enzymol* 197:24). *E. coli* phospholipids were metabolically labelled for 3 hours with [¹⁴C]-oleic acid (95% in *sn*-2 position). The bacteria were then autoclaved for
15 15min. to inactivate bacterial phospholipases, and washed. The viral sample (1-1000 pg) was added to the bacteria (10,000 cpm) in standard buffer (50 mM Tris-HCl, pH 8.0, and 10 mM CaCl₂) and incubated for 30min. at 37°C. The reaction was stopped by the addition of ice-cold 1% BSA. After centrifugation, 50% of the supernatant was counted.

The mixed micelles assay was also used to establish PLA₂ activity. The substrates (usually
20 from New England Nuclear/DuPont) were 6 µM of either L-3-phosphatidylcholine, 1,2-di[1-¹⁴C]oleoyl (specific activity 110 mCi/mmol), L-3-phosphatidylinositol L-1-stearoyl-2-[1-¹⁴C]arachidonyl (specific activity 48 mCi/mmol), or phosphatidylethanolamine L-1-palmitoyl, 2-[1-¹⁴C]arachidonyl (specific activity 54.6 mCi/mmol). This assay was based on that described by Manjunath *et al.*, (1994) *Biochem J.* 303:121, with the following
25 modifications: 1 mM Triton X-100 was used instead of deoxycholate; the reaction total volume was 50 µl and was stopped after 10 min by adding 80 µl chloroform/methanol (2:1) and 50 µl of saturated KCl solution. Separation was on silica gel thin layer chromatographic plates. Optimal separation of the phosphatidylcholine reaction products was obtained using

a solvent solution containing chloroform, methanol and water (65:35:4). For phosphatidylethanolamine and phosphatidylinositol, chloroform and methanol (87:13) were used as solvent. The separated products were quantified using a Molecular Dynamics PhosphorImager SI after drying. Bee venom PLA₂ was purchased from Sigma-Aldrich Co. 5 (Cat. No. P9279). Only the regression lines of activity vs. dilution with a correlation coefficient $r^2 > 0.98$ were used to calculate the amount of protein to hydrolyze 50% of the substrate in the mixed micelles assay. Activity was then expressed as the μmol phospholipid hydrolyzed during the 10min. assay per ng of enzyme.

Alternatively, the first-order rate constant, k , was calculated from reaction progress curves 10 using the integrated first-order rate equation $[P] = S_0(1 - e^{-kt})$, where P is product concentration and S is substrate concentration, and in which $k = (k_{\text{cat}}/K_M)_{\text{app}} E_0$, where E_0 is the total enzyme concentration. The value of $(k_{\text{cat}}/K_M)_{\text{app}}$ represents the catalytic efficiency of the enzyme.

MIMIC Assay:

The MIMIC assay is described in Payan *et al.*, (1997) *J. Virol. Methods* 65:299 and 15 Haberhausen *et al.*, (1998) *J. Clin. Microbiol.* 36:628. The primers used for MIMIC PCR were (positions 836-862 in NADL-2) 5'-AGTGGGTATCGCTACTAACCTACACTC [SEQ ID NO: 47] and (positions 1207-1181 in NADL-2) 5'-GATCTGTCATCATCCAGTCTTCTATGC [SEQ ID NO: 48]. The competing MIMIC amplicon contained a deletion from positions 863-903 and was cloned into a pBluescript 20 plasmid.

Immunofluorescence:

For immunofluorescence studies, the 3C9 monoclonal antibody (ATCC CRL-1745) was used as primary antibody (1 hour incubation). Goat anti-mouse IgG, labelled with fluorescein isothiocyanate was obtained from Sanofi Diagnostics Pasteur (Redmond, WA) and used as 25 secondary antibody (incubation 1 hour), following standard techniques.

Confocal Immunofluorescence:

The double labelling of internalized PPV with LAMP-2 was performed using a porcine polyclonal anti-PPV antibody and the AC17 mouse anti-canine LAMP-2 monoclonal

antibody (previously reported to recognize mink LAMP-2; Hariri *et al.*, (2000) *Mol. Biol. Cell* 11:255). Appropriate controls were performed to ensure that the antibody labelling was specific. *In situ* hybridization was performed using DIG-conjugated anti-VP1 up and anti-NS1 probes, and FITC-conjugated anti-DIG antibody sandwich labelling according to the
 5 manufacturer's instructions (Roche Molecular Biochemicals). Nuclear labelling was performed by the addition of 0.02 mg/ml propidium iodide to paraformaldehyde-fixed cells after labelling. Confocal images were acquired using the 63X PlanApochromat objective of a Leica TCS SP confocal microscope equipped with argon and krypton laser sources.

Virus Inactivation:

10 PPV, at a concentration of about 10^{10} GE/ml, was treated with β -propiolactone (BPL) at 0.1% for 8 hours at 18°C. BPL reacts preferentially with purines, with the main targets being N-1 at adenosine, N-3 at cytidine, and N-7 at guanosine. Depurination may be an important consequence (Brown *et al.*, (1974) *J. Virol.* 14:840; Drinkwater *et al.*, (1980) *Biochemistry* 19:5087). These conditions completely inactivated the virus. The viral capsid was not
 15 affected and remained fully reactive to the different monoclonal antibodies (obtained from ATCC).

EXAMPLE I: PROTEIN SEQUENCE COMPARISONS

The parvovirus genome is depicted in Figure 1. The right-half of the parvovirus genome contains a large ORF coding for a nested set of capsid proteins (VPs) from alternative in-
 20 frame initiation codons so that these VPs differ by their *N*-terminal extensions. Initiation codons are determined by alternative splicing (eg., VP1 and VP2 of porcine parvovirus (PPV) and human parvovirus B19) or a leaky-scanning mechanism (e.g. VP1-4 of *Galleria mellonella* densovirus, *GmDNV*). The VPs can be subdivided into 3 domains: common *C*-terminal domains required for capsid formation (60 copies per capsid); a small hinge-(like)
 25 domain; and unique *N*-terminal extensions. The capsid-forming domain consists of a β -barrel in which β -strands (black boxes: α A and α B are helices) are connected by loops (L1-4). These loops form most of the surface of the capsids, may contain allotropic determinants

(ats), and intertwine with neighboring subunits. Only the common domains are required to obtain stable capsids from expression vectors (Hernando *et al.*, (2000) *Virology* 267:299; Le Gall-Recule *et al.* (1996), *J Gen Virol* 77:2159; Miyamura *et al.*, (1994) *Proc Natl Acad Sci USA* 91:8507; Wong *et al.*, (1994) *J Virol* 68:4690; Brown *et al.*, (1994) *Virology* 198:477; 5 Martinez *et al.*, (1992) *Vaccine* 10:684). The known 3D-structure of 4 different parvoviruses, solved by X-ray crystallography, is restricted to these domains (Simpson *et al.*, (1998) *Structure* 6:1355; Agbandje-McKenna *et al.*, (1998) *Structure* 6:1369; Agbandje *et al.*, (1993) *Proteins* 16:155; Tsao *et al.*, (1991) *Science* 251:1456). The hinge-regions for the group of parvoviruses consisting of PPV, Minute Virus of mice (MVM), mink enteritis 10 parvovirus (MEV), canine parvovirus (CPV), and feline panleukemia parvovirus (FPV) contains a flexible polyglycine stretch that may enable the unique N-terminal extensions to be externalized through the 5-fold channel during maturation or early during infection (Cotmore *et al.*, (1999) *Virology* 254:169). The N-terminal extensions of B19 VPs, and probably of GmDNV, are permanently on the outside of the virion, and their hinge-like 15 regions have a high content of glycine, threonine and serine. About 100 amino acids upstream of the start of VP is a conserved domain (cd). The shaded boxes in Figure 1 represent the domains (V1ups) cloned in expression vectors (VP1up: PPV-VP1up amino acids 2-174 [SEQ ID NO: 50]; B19-VP1up amino acids 2-240 [SEQ ID NO: 52]; and GmDNV-VP1up amino acids 1-378 [SEQ ID NO: 54]).

20 Various protein sequences of the conserved domain (cd) region of the VP1up region of the largest capsid protein of 34 human, animal, and insect parvoviruses were compared by sequence alignment (Figure 2). The following parvovirus VP1ups were included in the sequence alignment: *Galleria mellonella* densovirus (Gm DNV) [SEQ ID NO: 1]; *Mythimna loreyi* densovirus (Ml DNV) [SEQ ID NO: 2]; *Junonia coenia* densovirus (Jc DNV) [SEQ 25 ID NO: 3]; *Pseudoplusia includens* densovirus (Pi DNV) [SEQ ID NO: 4]; *Diatraea saccharalis* densovirus (Ds DNV) [SEQ ID NO: 5]; *Culex pipiens* densovirus (Cp DNV) [SEQ ID NO: 6]; *Periplaneta fuliginos* densovirus (Pf DNV) [SEQ ID NO: 7]; *Acheta domesticus* densovirus (Ad DNV) [SEQ ID NO: 8]; *Casphalia extranea* densovirus (Ce DNV) [SEQ ID NO: 9]; *Bombyx mori* densovirus (Bm DNV) [SEQ ID NO: 10]; canine parvovirus 30 [SEQ ID NO: 11]; mink enteritis parvovirus [SEQ ID NO: 12]; mouse parvovirus 1 [SEQ

ID NO: 13]; feline panleukopenia parvovirus [SEQ ID NO: 14]; Minute Virus of Mice (MVM) [SEQ ID NO: 15]; Kilham rat parvovirus (K.Rat PV) [SEQ ID NO: 16]; porcine parvovirus [SEQ ID NO: 17]; Muscovy duck parvovirus (Mduck PV) [SEQ ID NO: 18]; goose parvovirus [SEQ ID NO: 19]; bovine parvovirus [SEQ ID NO: 20]; simian parvovirus
 5 [SEQ ID NO: 21]; chipmunk parvovirus [SEQ ID NO: 22]; Adeno-Associated Virus 2 (AAV2) [SEQ ID NO: 23]; Adeno-Associated Virus 3B (AAV3B) [SEQ ID NO: 24]; Adeno-Associated Virus 4 (AAV4) [SEQ ID NO: 25]; Adeno-Associated Virus 5 (AAV5) [SEQ ID NO: 26]; Adeno-Associated Virus 6 (AAV6) [SEQ ID NO: 27]; LuIII parvovirus (isolated from tissue culture) [SEQ ID NO: 28]; H1 parvovirus (isolated from tissue culture)
 10 [SEQ ID NO: 29]; human B19 parvovirus [SEQ ID NO: 30].

Representatives of sPLA₂s included in the alignment are: IA - *Naja naja* snake venom PLA₂ (amino acids 47 - 81 [SEQ ID NO: 31] and amino acids 115 - 126 [SEQ ID NO: 32]); IB - human pancreatic PLA₂ (amino acids 43 - 77 [SEQ ID NO: 33] and amino acids 116 - 127 [SEQ ID NO: 34]); IIA - human synovial fluid PLA₂ (amino acids 40 - 74 [SEQ ID NO: 35]
 15 and amino acids 106 - 117 [SEQ ID NO: 36]); IIB - gaboon viper snake venom PLA₂ (amino acids 18 - 52 [SEQ ID NO: 37] and amino acids 81 - 92 [SEQ ID NO: 38]); IIC - rat PLA₂ (amino acids 48 - 82 [SEQ ID NO: 39] and amino acids 118 - 129 [SEQ ID NO: 40]); III - Bee venom PLA₂ (amino acids 29 - 69 [SEQ ID NO: 41] and amino acids 87 - 98 [SEQ ID NO: 42]); V - human PLA₂ (amino acids 40 - 74 [SEQ ID NO: 43] and amino acids 106 -
 20 117 [SEQ ID NO: 44]); and X - human PLA₂ (amino acids 61 - 95 [SEQ ID NO: 45] and amino acids 128 - 139 [SEQ ID NO: 46]).

This analysis revealed the presence of a short stretch of high similarity, although in some cases the coding sequence was interrupted by an intron. Analysis of the 34 parvovirus sequences in this region also revealed a fully conserved HDXXY motif in all but four of the
 25 sequences (Aleutian Disease Virus and those of the Brevidensovirus genus). This HDXXY motif is also present in the catalytic site of secreted PLA₂s (sPLA₂s) (Figure 2).

In the sPLA₂ family (Dennis (1994) *J. Biol. Chem.* 269:13057-13060), the conserved H48 is assisted by D99 to polarize a catalytic water molecule that hydrolyzes the phospholipid ester. The conserved D49 and carbonyl oxygens from G30 and G32 bind a calcium ion involved

in the stabilization of the transition state. The conserved calcium-binding loop motif YXGXG is also present in VP1up, although not at the same sequence distance as in sPLA₂ (Fig. 2). Although sPLA₂s contain 5-8 disulfide bridges, which are a basis for their classification, these are absent in the parvovirus VP1up.

- 5 The comparison of VP1up sequences with those of sPLA₂ revealed major differences (in addition to disulfide bridges) that suggested vPLA₂ should be considered as a new group of the PLA₂ superfamily. The region connecting the HD helix and the D helix (Dijkstra *et al.*, (1981) *Nature* 289:604-606; White *et al.*, (1990) *Science* 250:1560-1563; Thunnissen *et al.*, (1990) *Nature* 347:689-691; Scott *et al.*, (1991) *Science* 254:1007-1010; Scott *et al.*, (1990)
- 10 *Science* 250:1563-1566) was found to be minimal in vPLA₂ (Fig.2). Although, the VP1up from the vertebrate B19 parvovirus showed some sequence identity (e.g., NPYTH) to the group IB pancreatic phospholipase, significant homology was also observed (Fig.2) between the vPLA₂ and group III PLA₂ (e.g., bee venom) at the amino-terminal and in the C-terminal helices which form the hydrophobic channel for the fatty acid chain (Scott *et al.*, (1991)
- 15 *Science* 254:1007-1010). The observed sequence homology would support the idea that vPLA₂ has a similar fold to the group III PLA₂ domain with connecting loop residues of the group I-II PLA₂ resulting in an overall different 3D structure.

- Searches of the databases Prosite PS00118 and PS00119, PRINTS PR00389, Pfam PF00068 and ProDom PD000303 using standard motifs used to recognize PLA₂ failed to identify
- 20 parvoviral phospholipase A₂ (pvPLA₂). Most probably this failure was due to the fact that these standard motifs rely heavily on the presence of cysteines which are necessary for disulfide bond formation and which are absent in pvPLA₂.

- Therefore, the conservation of the catalytic site and the Ca²⁺-binding loop was used to develop new motifs that allowed us to detect known and potential PLA₂s. Searches conducted
- 25 using the pattern [WY]-[CIVL]-G-x-[GP]-[GKNS]-x(8,14)-D-x(2)-[AC]-x(2)-H-D-x(2)-Y, and allowing for 2 mismatches using, e.g. the PATTINPROT program (PBIL (<http://npsa-pbil.ibcp.fr/>)), or Pattern and Profile Searches (ExPASy (<http://www.expasy.ch/>)), identified enzymatically active sPLA₂s from all sPLA₂ groups, albeit with high background. Refined,

more specific motifs recognized subsets of sPLA₂ representing specific sPLA₂ groups, with high specificity and sensitivity. In the above motif, the one-letter notation of amino acids is used, bracketed amino acids (eg [WY]) represent alternative amino acids at that position, x represents any amino acid, x(2) represents a stretch of 2 amino acids and x(8,14) a stretch of 5 between 8 and 14 amino acids.

For pvPLA₂, the specific motif is Y-x-G-P-G-x(12)-D-x(2)-A-x(2)-H-D-x(2)-Y. A search of the TrEMBL protein sequence database conducted using this motif detected all 102 vPLA₂ sequences, of which 5 had 1 mismatch. No additional pvPLA₂s were detected when 2 mismatches were allowed. Only when 3 or more mismatches were allowed were non-specific hits generated.

The general search patterns also permitted the detection of potential sPLA₂ and PLA₂ in other viruses. Virus infections have been reported to modify intracellular second-messenger pathways, lipid metabolism and arachidonic acid metabolism through pathways that normally are associated with PLA₂ (Abubakar *et al.* (1990) *Biochem. Biophys. Res. Commun.* **166**: 953; Shibutani *et al.*, (1997) *J. Clin. Invest.* **100**: 2054) thus indicating the importance of identifying potential PLA₂s in other viruses.

EXAMPLE II: PLA₂ ACTIVITY IN VP1up

The PLA₂ activity in the VP1up region of three divergent parvovirus genera was measured. Human B19, porcine parvovirus (PPV), and the insect parvovirus GmDENV were chosen as typical representatives to demonstrate the presence of viral PLA₂ (vPLA₂) and to identify its role in the viral cycle.

PLA₂ activity of VP1up was established using the *E. coli* radioactive assay and the mixed micelles assay, as described above, using sPLA₂s from snake venom, bee venom, and pig pancreas as controls. Although the *E. coli* radioactive assay is more sensitive than the mixed micelles assay, the modified mixed micelles assay was found to be particularly reproducible. PLA₂ activity in purified virus was only detected when high concentrations of PPV virions

($>5 \mu\text{g/ml}$) were assayed due to the fact that the PLA_2 domain resides predominantly inside the capsid at this stage. Alkali-denaturation and renaturation to disrupt the capsids and expose the VP1up increased the PLA_2 activity 50-100 fold to levels close to that of expressed VP1up. A heat-shock of 2min. at 70°C also released some VP1up as the PLA_2 activity increased about 20-50 times using this method. Polyclonal antibodies raised against PPV VP1up significantly reduced PLA_2 activity of VP1up exposed after dissociation of capsids or by heat shock, but did not affect the activity of bee venom PLA_2 .

With the mixed micelles assay, expressed VP1up of PPV had a 41-fold higher PLA_2 specific activity (relative molar amounts of enzyme required to hydrolyze 50% of the substrate) than bee venom s PLA_2 (which had a specific activity 1360 U/mg). B19 and GmDNV VP1up were 13.5 and 88.0 times less active, respectively, than bee venom PLA_2 , with detection thresholds in the low pg to ng range. The activity and pH optimum depended on the substrate. Although PPV and B19 v PLA_2 had similar activities at their respective pH optima (8.0 and 4.5) in the *E. coli* assay, B19 v PLA_2 had the same pH optimum as PPV v PLA_2 (pH 8.0) in the mixed micelles assay, but with a 550-fold lower activity.

The optimum Ca^{2+} concentration was similar to the extracellular (and some intracellular compartments) free Ca^{2+} concentration of about 1.2 mM (Evenas *et al.*, (1998) *Curr Opin Chem Biol* 2:293). A residual activity of 10% at $10 \mu\text{M}$ Ca^{2+} could be measured. Addition of EDTA or EGTA (chelator of Ca^{2+}) reduced activity of PPV and B19 v PLA_2 to below detectable levels in the mixed micelles assay (Figure 3). No activity was observed at the resting cell cytoplasmic Ca^{2+} concentration of $0.1 \mu\text{M}$, although this concentration may increase up to $10 \mu\text{M}$ in activated cells (Evenas *et al.*, (1998) *Curr Opin Chem Biol* 2:293) or infected cells (Hallett *et al.*, (1982) *Biochem J.* 206:671; Hartshorn *et al.*, (1988) *J. Immunol.* 141:1295; Irurzun *et al.*, (1995) *J Virol* 69:5142).

Reducing agents were not expected to influence the enzyme activity due to the absence of disulfide bridges. In fact, an increase in activity was observed in the presence of 2-mercaptoethanol (see Table 1).

**Table 1. PLA₂ activity (% of activities)
(parallel *E. coli* radioactive assay used unless noted otherwise)**

| | B19 | PPV |
|--------------------------------------------------------------------------------------------------|-----------------------|------------------------|
| pH optimum (<i>E. coli</i> assay) | 4.5 | 8 |
| pH optimum (mixed micelles assay) | 6 | 8 |
| 5 Ca ²⁺ | | |
| 1 μ M | 9.6 % | 8.6 % |
| 1 mM | 105.9 % | 68.0 % |
| 10 mM | 100 % | 100 % |
| 2-Mercaptoethanol | | |
| 10 10 mM | 148 % | 189 % |
| 100 mM | 154 % | 161 % |
| Monoclonal antibody to C-terminus of PPV VP1up (at about 0.5 μ g/50 μ l) ¹ | 100 % | 6.7 % |
| Inhibitors ² | | |
| 15 IC ₅₀ , manoalide, pH 8.0 | 3.8 \pm 0.4 μ M | 2.0 \pm 0.3 μ M |
| IC ₅₀ , oleyloxyethylphosphoryl choline, pH 8.0 | >20 μ M | 11.1 \pm 0.9 μ M |

¹Determined by use of a mixed micelles assay.

²The IC₅₀ is the inhibitory concentration leading to 50% inhibition.

The values of $(k_{cat}/K_M)_{app}$ for the PLA₂ activity of the expressed V1ups was also determined from the mixed micelle assays and are shown in Table 2.

Table 2. $(k_{cat}/K_M)_{app}$ Values

| Phospholipase A ₂ Source | $(k_{cat}/K_M)_{app} / M^{-1} s^{-1}$ |
|-------------------------------------|---------------------------------------|
| Bee venom | (1.09 \pm 0.14) $\times 10^5$ |
| Porcine PV V1up | (71.9 \pm 9.4) $\times 10^5$ |
| 25 B19 PV V1up | (2.5 \pm 0.2) $\times 10^4$ |
| GmDNV V1up | (0.4 \pm 0.03) $\times 10^4$ |
| BmDNV V1up | $\sim 0.5 \times 10^5$ |
| CeDNV V1up | $\sim 0.5 \times 10^5$ |
| AAV VP1up | $\sim 0.75 \times 10^5$ |

Extended regions of PPV VP1up, each containing the conserved pvPLA₂ domain, were also expressed and their PLA₂ activity was determined (Figure 3B). VP1up from M1 to S174 was used throughout this work and was assigned the relative specific activity of 1.0. Sequences within this expressed peptide but outside the PLA₂ motif (dark-grey in the figure) contributed to the activity as shown by the relative specific activities. The activities of these extended VP1up regions indicated that sequences outside the conserved domain had a large impact on the catalytic efficiency of the enzyme.

The PLA₂ activity of untreated virions (0.2 µg), virions after dissociation and after heat shock was also measured, using bee venom (18 ng) as a control. See Figure 3D. Samples in lanes 3, 5 and 7 were treated with anti-VP1up antibody (200 ng). This antibody reduced viral, but not bee venom, PLA₂ activity.

EXAMPLE III: SPECIFICITY OF PLA₂ ACTIVITY

The specificity of the vPLA₂ reaction was determined by thin layer chromatography (TLC) after hydrolysis of phosphatidylcholine substrate (PL) into fatty acid (FA) and lysophosphatidylcholine (LPL) by expressed vPLA₂ in the mixed micelles assay. PLA₂ activity was analyzed with a Molecular Dynamics PhosphorImager (%: fraction hydrolyzed in % = ((FA+LPL)/(FA+PL+LPL))x100). PLA₂s used in the assay were: PPV PLA₂ (0.6 ng), B19 PLA₂ (250 ng), GmDENV PLA₂ (350 ng) and bee venom PLA₂ (15 ng). Thioredoxin (2,000 ng) was used as a negative control. The results are shown in Figure 3A. The 3C9 monoclonal antibody, which binds to the C-terminus of PPV-VP1up, reduced PPV PLA₂ activity of VP1up to 18% of the original activity after adding 200 ng of antibody (lane 2), but not that of B19 (lane 4). Adding EGTA to a final concentration of 5 mM abolished PLA₂ activity (lane 5).

Following hydrolysis, TLC showed that the radioactive label, if present only in the *sn*-2 oleic acid, moved from the phospholipids to the fatty acid (free oleic acid) spots, and, when present

at both *sn*-1 and *sn*-2 positions, to both the lysophosphatidyl and fatty acid spots (Figure 3A), thus confirming the specificity of the PLA₂ activity.

The impact of Ca²⁺ concentration, pH and different substrates on the activity of pvPLA₂ was also investigated. Since 1 mM Ca²⁺, 50 mM Tris-HCl, pH 8.0, and phosphatidylcholine (PC) were used in standard assays, these were set at 100% relative activity. The results of these assays are shown in Figure 3C. PPV can be seen to efficiently hydrolyse both phosphatidylethanolamine (PE) and phosphatidylinositol (PI), the alternative substrates tested.

EXAMPLE IV: INHIBITING PLA₂ ACTIVITY

PLA₂ inhibitors manoalide (MA) and oleyloxyethylphosphorylcholine (OP) inhibited the VP1up PLA₂ activity in the *E. coli* assay. The MA concentration leading to 50% inhibition of PLA₂ was 3.8±0.4 for B19 and 2.0±0.3 µM for PPV. Using OP, >20 and 11.1±0.9 µM, respectively, was required to achieve the same degree of inhibition.

EXAMPLE V: 3-D STRUCTURE OF vPLA₂

The 3D-structures of a large number of group I/II sPLA₂ and group III sPLA₂ have been solved (e.g., Renetseder *et al.*, (1985) *J. Biol. Chem.* 260:11627; Yuan and Tsai (1999) *Biochim. Biophys. Acta.* 1441:215; Dijkstra *et al.*, (1981) *Nature* 289:604); Sekar and Sundaralingam (1999) *Acta. Crystallogr. D. Biol. Crystallogr.* 55:46; van den Berg *et al.*, (1995) *Embo J.* 14:4123; Dijkstra *et al.*, (1978) *J. Mol. Biol.* 124:53). This permitted the prediction of the B19/PPV vPLA₂ structure, which then served to select potentially critical amino acids for site-directed mutagenesis (Figure 4). This sequence and structure alignment indicated that neither the position numbering of group I/II nor that of group III could be maintained and resulted in an adapted PPV numbering for vPLA₂.

EXAMPLE VI: SITE-DIRECTED MUTAGENESIS TO CREATE PLA₂ MUTANT PARVOVIRUSES

The 3-D structure of vPLA₂ was used to identify potentially critical amino acids in this enzyme. From this structure, residues 21 (proline), 35 (aspartic acid), 41 (histidine), 42
5 (aspartic acid), 63 (aspartic acid) and 88 (lysine) were selected for site-directed mutagenesis experiments.

Both the PPV-VP1up and the virus itself were submitted to site-directed mutagenesis (Weiner *et al.*, (1994) *Gene* 151:119; Kuipers *et al.*, (1990) *Prot. Eng.* 3:599). The mutants of the infectious clone of PPV (NADL-2 strain in pUC19 (Bergeron, *et al.*,
10 (1996) *J Virol* 70:2508; Maxwell (1993) *J. Gen. Virol.* 74:1175; Brownstein, *et al.*, (1992) *J Virol* 66:3118; Gardiner and Tattersall (1988) *J Virol* 62:1713; Corsini, *et al.*, (1996) *Adv Virus Res* 47:303; Berns (1990) *Microbiol Rev* 54:316)) were transfected into PT cells and large amounts were produced (despite their often low infectivity). Particular attention was devoted to detect possible revertants by monitoring the infectivity in two
15 successive infection assays. The concentration of full virus particles, expressed as genome equivalents or GE per µl, was determined by a MIMIC assay. The infectivity of the suspensions was measured with the fluorescent-focus assay. The number of fluorescent nuclei in a well was counted using immunofluorescence 20 hours after infection (before fluorescent nuclei from a secondary infection could appear). The relative
20 infectivity of the mutants was determined from the number of full particles, as established by GE in the MIMIC assay, required to obtain one fluorescent focus unit (ffu), after comparing it to the wild-type (wt) virus. Dilutions were used that gave at least 25 ffu (relative error <0.2). For wt PPV, 232 full particles were required per ffu, and this value represents 100%. The value for the relative specific infectivity of each mutant represents
25 the result of five independent assays.

The PLA₂ activity of the wt and mutant VP1ups was determined as shown in Table 3. Hydrolysis of the substrate was measured with a PhosphorImager. Different amounts of expressed pvPLA2 were assayed and the degree of hydrolysis was measured. In this assay,

0.46 ng wt PPV pvPLA₂ fusion protein was required to achieve 50% hydrolysis. The relative amount of each mutant required to obtain the same degree of hydrolysis yielded the relative specific activities. Only the regression lines of activity vs dilution with a correlation coefficient $r^2 > 0.98$ were used to calculate the amount of protein to hydrolyze
5 50% of the substrate in the mixed micelles assay.

Both the enzyme activity and viral infectivity decreased significantly when amino acids in the catalytic site (H41 and D42) were mutated (Table 3), whereas back-mutations restored the original infectivity/activity (the back-mutations served as a control to confirm the integrity of rest of genome). Among potential aspartic acids corresponding to D99 in
10 sPLA₂, D63 was fully conserved in vPLA₂ (Figure 2), and mutation of this residue was found to decrease strongly both infectivity and activity, indicating that the relative position of the predicted α -helices would be correct (Figure 4).

Site-directed mutagenesis of the aspartic acid residue at position 35, one of the very few
15 amino acids that is conserved among all sPLA₂ and vPLA₂ to glutamic acid (D35E) or asparagine (D35N), had an impact that was comparable to the D63E or D63N mutations. This amino acid may be critical for the positioning of the Ca²⁺-loop by 3 hydrogen bonds to this loop.

A surprising observation was the conservation of P21 among the vPLA₂ but not among
20 the sPLA₂. When P21 was mutated to amino acids that occur at that position in sPLA₂ (R, W, L), both the infectivity and the activity were strongly reduced. It is not known whether mutations to P21 in sPLA₂ would increase their activities. This residue has a versatile role (Kuipers *et al.*, (1990) *Protein Eng* 3:599.) such as involvement in binding and orienting monomeric substrate, binding of the enzyme to micellar substrates and
25 possibly shielding the catalytic site from excess water. P21 may be required in the case of vPLA₂ to compensate for the 3-amino acid insertion between the Ca²⁺-binding loop and the α -helix containing H41.

K88 is conserved among parvoviruses, and in bee venom (K85) where it is involved in receptor binding (Nicolas *et al.*, (1997) *J. Biol. Chem.* 272:7173). The potential role of this residue as a co-receptor in parvoviruses could not be established since conservative mutagenesis (K88R) in VP1up already decreased activity strongly (no receptor involved).

- 5 Nevertheless, the presence of this conserved homologue of the bee venom PLA₂ receptor-binding helix is intriguing.

Table 3. Relative Specific Activity and Infectivity of pv PLA₂ Mutants

| 10 | pvPLA ₂ mutants (PPV) | Position in group I/II | Position in group III | Relative specific activity (%) | Relative specific infectivity (%) |
|----|----------------------------------------|---------------------------|--------------------------|-----------------------------------|--------------------------------------|
| | wt | | | 100 | 100 |
| | P21L | 31 | 11 | 7.5 | 0.62 |
| | P21R | 31 | 11 | 0.07 | 0.0012 |
| | P21W | 31 | 11 | 4.1 | 0.055 |
| 15 | D35E | 42 | 28 | 0.27 | 0.071 |
| | D35N | 42 | 28 | 0.02 | 0.0019 |
| | H41A | 48 | 35 | <0.007 | 0.0011 |
| | D42N | 49 | 36 | 0.04 | 0.0012 |
| 20 | HD41/42A N | 48/49 | 35/36 | <0.005 | 0.0011 |
| | D63A | 99 | 64 | <0.003 | 0.0003 |
| | D63N | 99 | 64 | 0.05 | 0.0007 |
| | K88R | - | 85 | 0.01 | 0.0035 |

EXAMPLE VII: IMMUNOFLUORESCENCE OF INFECTED CELLS

The observation that, upon transfection, wild-type and mutant infectious clones are both effective in producing virions indicated that vPLA₂ is required at some stage prior to replication, i.e. during entry. In order to assess the critical step(s), cells were infected with wt or mutant virus (P21L and P21W) at a concentration of 0.5 µg/ml and tracked by immunofluorescence. Virus was adsorbed to nonconfluent cells for 5 hours at 4°C to ensure synchronized entry when the cold medium was replaced by medium at 37°C (time-point 0 hours).

The following stages could be distinguished for wt by immunofluorescence (Figure 5): 0 hrs: strong, patchy staining on the cellular membrane; 4 hrs: virus entry with cytoplasmic, granular and perinuclear staining; 8 hrs: first nuclear staining; and 12 and 18 hrs: strong nuclear staining. Prior addition of lysosomotropic NH₄Cl reduced infection in a dose-dependent fashion (10, 20, and 50 mM resulted in 20-, 100- and 100,000-fold reduction in infectivity). At 20 mM NH₄Cl, the first nuclear staining was delayed to 18 hrs. Although the entry was fast (5-30 min), a decrease in infectivity could be obtained up to about 4 hrs post-infection by adding NH₄Cl or neutralizing antibodies suggesting a long stay in vesicular compartments. These results are consistent with the observation that parvovirus enters the cell via receptor-mediated endocytosis (Parker and Parrish (2000) *J. Virol.* 74:1919; Basak and Turner (1992) *Virology* 186:368; Vihinen-Ranta *et al.*, (1998) *J. Virol.* 72:802). It has been observed (Parker and Parrish (2000) *J. Virol.* 74:1919) that canine parvovirus (CPV) co-localizes with transferrin in perinuclear endosomes, suggesting that parvovirus infection is complex and involves multiple vesicular components. Biotinylation of virus prior to infection and detection by streptavidin-fluorescein, which only detects incoming virus, showed perinuclear staining after 4, 8, 12, 18 hrs but no nuclear staining. In contrast, immunofluorescence showed nuclear staining from about 8 hrs. It also indicated that infectivity, as measured by ffu, was reduced to about 50% by biotinylation. This perinuclear accumulation was not due to biotinylation since the same localization was obtained after inactivation of the virus without affecting the capsid (see Methods and Materials), infection, or subsequent immunofluorescence at 4, 8, 12, and 18 hrs. Although this suggests that only the viral genome is delivered from the perinuclear zone, perhaps from vesicles, into the nucleus, it remains possible that a

few virus particles entering the nucleus, which are below the detection level, are responsible for the infection.

Several differences were observed between the mutants and the wt virus. In contrast to the wt virus, the mutants showed a rather diffuse staining early during entry. Like the wt, the mutants became perinuclear after 4 hrs and could not be distinguished at this point from wt. Mutants, however, remained perinuclear for long periods (at least 18 hrs) whereas wt virus-infected cells showed a strong nuclear staining starting after 8 hrs. Occasionally, an infected nucleus was obtained after 18 hrs with the mutants. NH_4Cl exacerbated the effect of the mutations. Although the incoming virus still became perinuclear at about 4 hrs post-infection, it remained there without leading to infection. Single-stranded parvoviral genomes entering the nucleus are normally converted into double-stranded DNA by cellular DNA polymerase (P. Tattersall and S.F. Cotmore in P. Tijssen, ed., *Handbook of Parvoviruses*, vol. 1 (Boca Raton: CRC Press, 1990) 123). This dsDNA would yield productive infections as was shown with mutant or wt infectious clones. Taken together, these results indicate that PLA_2 mutants experience difficulties in transferring their genome from the perinuclear localization into the nucleus.

EXAMPLE VIII. CONFOCAL IMMUNOFLUORESCENCE OF INFECTED CELLS

The binding and entry of ^{35}S -labeled PPV were studied by confocal immunofluorescence using 4×10^5 cells and 5×10^9 wild type virus particles. Equivalent amounts of mutant viruses were used for the binding studies, and twice as much for the entry studies. Experiments were performed in quadruplicate. The results are shown in Figure 6A (binding) and 6B (entry). Viruses that were either bound to cells or had entered into cells were measured by a liquid scintillation method after performing washing or antibody-stripping procedures, respectively. pvPLA_2 mutations had no effect on virus binding to the cells or entry. Both binding and entry of virus were time- and dose-dependent (not shown).

Figure 6C shows the relative radioactivity from experiments in which 10^6 cpm ^{35}S -labeled PPV particles (wt or mutant) were added to 2×10^7 cells in Petri dishes and about 20% was recovered in the cells. After cellular fractionation, the distribution of virus over cytosolic and nuclear fractions was measured by liquid scintillation. Relative radioactivity was calculated as follows: [nuclear fraction-bound activity]/[cellular fraction-bound activity]. Standard errors were calculated from three experiments.

In addition, co-localization of PPV and LAMP-2 was studied by confocal microscopy. Anti-PPV antibodies were labeled with FITC, giving green fluorescence, and anti-LAMP-2 antibodies with Texas Red, giving red fluorescence. Co-localization of PPV and LAMP-2 resulted in yellow staining. The results are shown in Figure 6D. Both wt and mutant viruses, as well as capsids lacking PLA₂, showed extensive co-localization with LAMP-2 in the late endosomes/lysosomes.

Taken together the above results indicate that parvoviruses do not require vPLA₂ activity for binding to the cell surface or initial stages of entry.

15 *EXAMPLE IX. IN SITU HYBRIDIZATION OF INFECTED CELLS*

Figure 7 shows the distribution of viral DNA in infected cells. Incoming viral DNA was detected in the cytoplasm by *in situ* hybridization 4 (A) and 8 (B) hours post-infection (p.i.), whereas replicating DNA in the nucleus was found at 12 hours p.i. (C). At 12 hours p.i., no DNA could be detected in the nucleus in the case of the HD (D) and P21^{PW} mutants.

20 Aphidicoline, a DNA polymerase inhibitor used at 2 $\mu\text{g/ml}$, also prevented the appearance of viral DNA in the nucleus (F). The sPLA₂ inhibitors tetracain (TC, at 100 μM), and oleyloxyethyl phosphorylcholine (OP, at 20 μM), also reduced the number of nuclei containing viral DNA (G).

EXAMPLE X. IDENTIFICATION OF VIRAL VLA₂-BINDING PEPTIDES

Four peptides that specifically bind to the PLA₂-containing VP1up of PPV using a phage display library. The Ph.D.TM Phage Display Peptide Library kit was obtained from New England Biolabs and used according to the manufacturer's instructions, with the exception that Linbro/Titertek (Cat. No. 76-231-05) plastic plates were used in the panning procedure
5 . to avoid high background.

The sequences peptides displayed on the phage that specifically bound to PLA₂ (p6, p7, p8 and p10) are shown in Figure 13. The nucleic acid sequences encoding these 12 amino acid peptides were subsequently cloned downstream of thioredoxin in the pBADTBX vector using the *Bgl*II and *Xba*I cloning sites, and expressed as fusion proteins.

- 10 The fusion proteins were used in ELISA assays to determine their ability to bind the PLA₂-containing VP1up of PPV. The titers obtained in ELISA with these fusion proteins and using anti-thioredoxin antibodies as secondary antibody were all, except for p10, about 3000. P10 was about 5x weaker.

The invention being thus described, it will be obvious that the same may be varied in many
15 ways. Such variations are not to be regarded as a departure from the spirit and scope of the invention, and all such modifications as would be obvious to one skilled in the art are intended to be included within the scope of the following claims.

THE EMBODIMENTS OF THE INVENTION IN WHICH AN EXCLUSIVE PROPERTY OR PRIVILEGE IS CLAIMED ARE DEFINED AS FOLLOWS:

1. An isolated viral polypeptide that has phospholipase A₂ activity.
2. The isolated viral polypeptide according to Claim 1, wherein said polypeptide comprises an amino acid motif:
 [WY]-[CIVL]-G-x-[GP]-[GKNS]-x(8,14)-D-x(2)-[AC]-x(2)-H-D-x(2)-Y
 wherein:
 □ indicates the presence of one of the enclosed amino acids at this position;
 x is any amino acid;
 x(2) represents a stretch of 2 amino acids;
 x(8,14) represents a stretch of between 8 and 14 amino acids.
3. The isolated polypeptide according to Claim 2 wherein said amino acid motif is:
 Y-x-G-P-G-x(12)-D-x(2)-A-x(2)-H-D-x(2)-Y
 wherein:
 x(12) represents a stretch of 12 amino acids; and
 wherein one of the specified amino acids Y, G, P, G, D, A, D, or Y may be exchanged singly for any other amino acid.
4. The polypeptide according to any one of Claims 1 - 3 wherein said viral polypeptide is a parvoviral polypeptide.
5. The polypeptide according to any one of Claims 1 - 3 wherein said parvoviral polypeptide is derived from *Galleria mellonella* densovirus, *Mythimna loreyi* densovirus, *Junonia coenia* densovirus, *Pseudoplusia includens* densovirus, *Diatraea saccharalis* densovirus, *Culex pipiens* densovirus, *Periplaneta fuliginos* densovirus, *Acheta domesticus* densovirus, *Casphalia extranea* densovirus, *Bombyx mori* densovirus, canine parvovirus, mink enteritis parvovirus, mouse parvovirus 1, feline panleukopenia parvovirus, Minute Virus of Mice, Kilham rat parvovirus, porcine parvovirus, Muscovy duck

- parvovirus, goose parvovirus, bovine parvovirus, simian parvovirus, chipmunk parvovirus, Adeno-Associated Virus 2, Adeno-Associated Virus 3B, Adeno-Associated Virus 4, Adeno-Associated Virus 5, Adeno-Associated Virus 6, LuIII parvovirus, H1 parvovirus, or human B19 parvovirus.
6. The polypeptide according to Claim 5 wherein the parvovirus is human B19, porcine parvovirus, *Galleria mellonella* densovirus, *Casphalia extranea* densovirus, *Bombyx mori* densovirus or Adeno-Associated Virus.
 7. An isolated viral polypeptide comprising an amino acid sequence as set out in SEQ ID NO: 55; SEQ ID NO: 57 or SEQ ID NO: 59.
 8. An isolated viral polypeptide that has phospholipase A₂ activity and comprising an amino acid sequence as set out in SEQ ID NO:49; SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:55, SEQ ID NO:57 or SEQ ID NO:59.
 9. The polypeptide according to any one of Claims 1- 8 wherein said polypeptide is chemically synthesized.
 10. The polypeptide according to any one of Claims 1-9 wherein said polypeptide fused to a heterologous polypeptide.
 11. A polypeptide that is an inactive variant of a viral polypeptides wherein said viral polypeptide comprises an amino acid motif:
 [WY]-[CIVL]-G-x-[GP]-[GKNS]-x(8,14)-D-x(2)-[AC]-x(2)-H-D-x(2)-Y
 wherein:
 [] indicates the presence of one of the enclosed amino acids at this position;
 x is any amino acid;
 x(2) represents a stretch of 2 amino acids;
 x(8,14) represents a stretch of between 8 and 14 amino acids.

12. An isolated polynucleotide encoding a viral polypeptide that has phospholipase A₂ activity.
13. The isolated polynucleotide according to Claim 13, wherein said polypeptide comprises an amino acid motif:
[WY]-[CIVL]-G-x-[GP]-[GKNS]-x(8,14)-D-x(2)-[AC]-x(2)-H-D-x(2)-Y
wherein:
[] indicates the presence of one of the enclosed amino acids at this position;
x is any amino acid;
x(2) represents a stretch of 2 amino acids;
x(8,14) represents a stretch of between 8 and 14 amino acids.
14. The isolated polynucleotide according to Claim 14 wherein said amino acid motif is:
Y-x-G-P-G-x(12)-D-x(2)-A-x(2)-H-D-x(2)-Y
wherein:
x(12) represents a stretch of 12 amino acids; and
wherein one of the specified amino acids Y, G, P, G, D, A, D, or Y may be exchanged singly for any other amino acid.
15. The isolated polynucleotide according to any one of Claims 12 - 14, wherein said polynucleotide is DNA, cDNA, RNA or genomic DNA.
16. The isolated polynucleotide according to anyone of Claims 12 - 15, wherein said polynucleotide is double stranded or single stranded.
17. The isolated polynucleotide according to any one of Claims 12 - 16, wherein said polynucleotide is fused to a heterologous polynucleotide.
18. The isolated polypeptide according to Claim 17 wherein said heterologous polynucleotide encodes a heterologous polypeptide.

19. An isolated polynucleotide comprising a nucleic acid sequence as set out in SEQ ID NO:56; SEQ ID NO:58 or SEQ ID NO:60.
20. An isolated polynucleotide encoding a viral polypeptide that has phospholipase A₂ activity and comprising a nucleic acid sequence as set out in SEQ ID NO:50; SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:58 or SEQ ID NO:60.
21. An antisense oligonucleotide complementary to the polynucleotide according to any one of Claims 12 - 16, 19 or 20.
22. Use of the antisense oligonucleotide according to Claim 21 to inhibit viral replication and / or infectivity in a subject.
23. A vector comprising the isolated polynucleotide according to any one of Claims 12 - 20.
24. The vector according to Claim 23 wherein said polynucleotide is operatively linked to one or more regulatory sequences.
25. A method of producing a host cell comprising genetically engineering cells with the vector according to Claim 23 or 24.
26. The host cell produced according to Claim 25.
27. A host cell comprising the vector according to Claim 23 or 24.
28. A host cell comprising the isolated polynucleotide according to any one of Claims 12 - 20.

29. A method of producing a recombinant virus comprising genetically engineering viral particles with the polynucleotide, or fragment thereof, according to Claims 12 - 16, 19 or 20, or with the vector according to Claim 23 or 24.
30. The recombinant virus produced according to Claim 29.
31. An antibody that specifically binds to the polypeptide of any one of Claims 1-9.
32. The antibody according to Claim 17 or 18, wherein said antibody is polyclonal.
33. The antibody according to Claim 17 or 18, wherein said antibody is monoclonal.
34. Use of a polypeptide according to any one of Claims 1-9 as an *in vitro* standard for measuring phospholipase A₂ activity.
35. A method of screening compounds in order to identify an agent that inhibits viral phospholipase A₂ activity, comprising the steps of:
 - i) incubating a viral phospholipase A₂ and a phospholipid substrate with a candidate agent under conditions whereby, but for the presence of the agent said viral phospholipase would hydrolyse said phospholipid, and
 - ii) measuring the inhibition of viral phospholipase A₂ activity.
36. An anti-viral agent that selectively inhibits the activity of a viral phospholipase isolated according to the method of Claim 35.
37. The anti-viral agent according to Claim 36, wherein said anti-viral agent is capable of producing a detectable decrease in infectivity and / or replication of a virus.

38. The anti-viral agent according to Claim 36 or 37, wherein said anti-viral agent is an antisense oligonucleotide that binds specifically to mRNA encoding said viral phospholipase.
39. The anti-viral agent according to Claim 38 wherein said antisense oligonucleotide is about 8 to about 100 nucleobases.
40. The anti-viral agent according to Claim 39 wherein said antisense oligonucleotide is about 8 to about 50 nucleobases.
41. The anti-viral agent according to Claim 40 wherein said antisense oligonucleotide is about 15 to about 30 nucleobases.
42. The anti-viral agent according to Claim 36 or 37, wherein said anti-viral agent is an antibody wherein said antibody is monoclonal or polyclonal.
43. A method of identifying an anti-viral agent that selectively inhibits the activity of a viral phospholipase A₂, comprising:
 - i) selecting a candidate molecule on the basis of the capacity of the candidate molecule to inhibit viral phospholipase A₂ in an *in vitro* assay employing a predetermined amount of a standardized preparation of viral phospholipase A₂;
 - ii) treating a cultured host cell population with the candidate molecule;
 - iii) exposing the host cell population to a virus;
 - iv) determining the reduction in the infectivity and / or replication of the virus;
 - v) determining the growth of the host cell population; and
 - vi) comparing the growth to the growth of an untreated cultured host cell population that has been exposed to a virus;wherein a reduction in the infectivity and / or replication of the virus and increased growth of the host cell population in comparison to the untreated

host cell population is indicative of selective inhibition of the viral phospholipase A₂.

44. An anti-viral agent that selectively inhibits the activity of a viral phospholipase isolated according to the method of Claim 43.
45. The anti-viral agent according to Claim 44, wherein said anti-viral agent is capable of producing a detectable decrease in infectivity and / or replication of a virus *in vivo*.
46. The anti-viral agent according to Claim 44 or 45, wherein said anti-viral agent is an antisense oligonucleotide.
47. The anti-viral agent according to Claim 44 or 45, wherein said anti-viral agent is a monoclonal antibody.
48. Use of the anti-viral agent according to any one of Claims 36 - 42 or 44 - 47 to decrease the infectivity and / or replication of a virus in a subject.
49. The use according to Claim 48 wherein said virus is a parvovirus.
50. Use of the anti-viral agent according to any one of Claims 36 - 42 or 44 - 47 to inhibit a viral phospholipase A₂ in a subject in need of such therapy.
51. The use according to Claim 50 wherein said subject has a viral-associated disease or disorder.
52. The use according to Claim 51 wherein said viral-associated disease is associated with a parvovirus.

53. The use according to Claim 52 wherein said viral-associated disease is rheumatoid arthritis, systemic erythematosus, adult-onset Still's disease or polyarthritis.
54. Use of the anti-viral agent according to any one of Claims 36 - 42 or 44 - 47 to manufacture a medicament to decrease the infectivity and / or replication of a virus in a subject.
55. Use of the anti-viral agent according to any one of Claims 36 - 42 or 44 - 47 to manufacture a medicament to treat a viral-associated disease.
56. Use of the antisense oligonucleotide according to any one of Claims 38 - 41 to manufacture a medicament to decrease the infectivity and / or replication of a virus in a subject.
57. Use of the antisense oligonucleotide according to any one of Claims 38 - 41 to manufacture a medicament to treat a viral-associated disease.
58. Use of the isolated polynucleotide according to any one of Claims 12 - 20 for gene therapy.
59. Use of the isolated polynucleotide according to any one of Claims 12 - 20 to improve a gene therapy vector.
60. The use according to Claim 59 wherein said gene therapy vector is based on an Adeno-Associated virus.
61. A peptide that specifically binds a viral phospholipase A₂.
62. The peptide according to Claim 61 wherein said peptide inhibits the viral phospholipase A₂.




Figure 1

BEST AVAILABLE COPY

| | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
|--------------|-----|---------|----|----|----|---|-----|-----|---|---|----|---|---|---|---|---|---|---|---|---|---|---|---|---|-------|-----|---|-----|---|---|---|---|---|---|-------|-----|----------|-------|-------------|------------|-----|------------|------------|
| Gm DNV | 177 | ITVPGKX | GP | GN | S | N | --- | RQ | Q | T | NQ | I | D | E | A | H | D | E | X | D | K | V | K | T | --- | SQ | E | S | R | A | D | N | T | E | N | K | --- | 230 | (AAA66966) | | | | |
| M7 DNV | 177 | ITVPGKX | GP | GN | S | N | --- | RQ | Q | T | NQ | I | D | E | A | H | D | E | X | D | K | A | K | T | --- | SQ | E | S | E | A | D | N | T | E | N | K | --- | 230 | (Q90053) | | | | |
| Jc DNV | 177 | ITVPGKX | GP | GN | S | N | --- | RQ | Q | T | NQ | I | D | E | A | H | D | E | X | D | K | A | K | T | --- | SQ | E | S | Q | A | D | N | T | E | N | K | --- | 230 | (AAC18002) | | | | |
| Pj DNV | 177 | ITVPGKX | GP | GN | S | N | --- | RQ | Q | T | NQ | I | D | E | A | H | D | E | X | D | K | A | K | T | --- | SQ | E | S | D | A | D | N | T | E | N | K | --- | 230 | (AAC18002) | | | | |
| Ds DNV | 177 | ITVPGKX | GP | GN | S | N | --- | RQ | Q | T | NQ | I | D | E | A | H | D | E | X | D | K | A | K | T | --- | SQ | E | S | D | A | D | N | T | E | N | K | --- | 230 | (AAC18002) | | | | |
| Cp DNV | 140 | VPAPV | AG | GN | S | N | --- | RQ | P | A | Y | D | L | V | E | S | A | Q | I | T | D | K | A | K | S | --- | P | E | D | I | H | K | D | R | O | T | E | --- | 193 | (AAF04300) | | | |
| Pf DNV | 149 | ITYEFH | Y | GP | GN | P | D | --- | N | N | E | V | D | R | D | A | I | A | E | H | D | C | A | N | A | K | S | --- | S | T | D | V | I | N | D | K | A | D | H | --- | 202 | (AAF04300) | |
| Ad DNV | 178 | AVLGGT | D | GP | GN | P | D | --- | K | P | A | R | S | E | T | D | I | A | E | H | D | C | A | N | A | K | S | --- | S | T | D | V | I | N | D | K | A | D | H | --- | 234 | (AF375296) | |
| Ce DNV | 4 | HFHYHN | Y | GP | GN | F | --- | K | Q | V | D | E | D | I | A | E | H | D | C | A | N | A | K | S | --- | S | T | D | V | I | N | D | K | A | D | H | --- | 62 | (AF375296) | | | | |
| Bm DNV | 4 | HFHYHN | Y | GP | GN | F | --- | K | Q | V | D | E | D | I | A | E | H | D | C | A | N | A | K | S | --- | S | T | D | V | I | N | D | K | A | D | H | --- | 62 | (AY033435) | | | | |
| Canine PV | 33 | VPBGY | Y | GP | GN | S | D | --- | Q | E | R | T | N | P | S | D | A | A | E | H | D | E | A | A | Y | L | R | S | G | K | N | P | Y | L | F | S | P | A | D | Q | --- | 91 | (VCPVCP) |
| Mink PV | 7 | VPBGY | Y | GP | GN | S | D | --- | Q | E | R | T | N | P | S | D | A | A | E | H | D | E | A | A | Y | L | R | S | G | K | N | P | Y | L | F | S | P | A | D | Q | --- | 65 | (VCPVME) |
| Mouse1 PV | 7 | VPBGY | Y | GP | GN | S | D | --- | Q | E | R | T | N | P | S | D | A | A | E | H | D | E | A | A | Y | L | R | S | G | K | N | P | Y | L | F | S | P | A | D | Q | --- | 65 | (AAA61406) |
| Feline PV | 12 | VPBGY | Y | GP | GN | S | D | --- | Q | E | R | T | N | P | S | D | A | A | E | H | D | E | A | A | Y | L | R | S | G | K | N | P | Y | L | F | S | P | A | D | Q | --- | 70 | (AAC37928) |
| MVM PV | 1 | MVPPGY | Y | GP | GN | S | D | --- | Q | E | R | T | N | P | S | D | A | A | E | H | D | E | A | A | Y | L | R | S | G | K | N | P | Y | L | F | S | P | A | D | Q | --- | 59 | (VCPVIM) |
| LuIII PV | 12 | WVPPGY | Y | GP | GN | S | D | --- | Q | E | R | T | N | P | S | D | A | A | E | H | D | E | A | A | Y | L | R | S | G | K | N | P | Y | L | F | S | P | A | D | Q | --- | 70 | (M81888) |
| H1 PV | 12 | WVPPGY | Y | GP | GN | S | D | --- | Q | E | R | T | N | P | S | D | A | A | E | H | D | E | A | A | Y | L | R | S | G | K | N | P | Y | L | F | S | P | A | D | Q | --- | 70 | (P03136) |
| K.Rat PV | 12 | CVPPGY | Y | GP | GN | S | D | --- | Q | E | R | T | N | P | S | D | A | A | E | H | D | E | A | A | Y | L | R | S | G | K | N | P | Y | L | F | S | P | A | D | Q | --- | 70 | (AAB38327) |
| porcine PV | 11 | LTLP | Y | GP | GN | S | D | --- | Q | E | R | T | N | P | S | D | A | A | E | H | D | E | A | A | Y | L | R | S | G | K | N | P | Y | L | F | S | P | A | D | Q | --- | 69 | (VCPVNA) |
| MDuck PV | 53 | FVLP | Y | GP | GN | S | D | --- | K | P | V | N | K | A | D | S | V | A | L | H | D | E | A | A | Y | L | R | S | G | K | N | P | Y | L | F | S | P | A | D | Q | --- | 111 | (CAA52984) |
| Goose PV | 53 | FVLP | Y | GP | GN | S | D | --- | K | P | V | N | K | A | D | S | V | A | L | H | D | E | A | A | Y | L | R | S | G | K | N | P | Y | L | F | S | P | A | D | Q | --- | 111 | (AAA3230) |
| AAV2 PV | 45 | FVLP | Y | GP | GN | S | D | --- | K | P | V | N | K | A | D | S | V | A | L | H | D | E | A | A | Y | L | R | S | G | K | N | P | Y | L | F | S | P | A | D | Q | --- | 103 | (AAC03780) |
| AAV3B PV | 45 | FVLP | Y | GP | GN | S | D | --- | K | P | V | N | K | A | D | S | V | A | L | H | D | E | A | A | Y | L | R | S | G | K | N | P | Y | L | F | S | P | A | D | Q | --- | 103 | (AAB95452) |
| AAV4 PV | 44 | FVLP | Y | GP | GN | S | D | --- | K | P | V | N | K | A | D | S | V | A | L | H | D | E | A | A | Y | L | R | S | G | K | N | P | Y | L | F | S | P | A | D | Q | --- | 102 | (AAC58045) |
| AAV5 PV | 44 | FVLP | Y | GP | GN | S | D | --- | K | P | V | N | K | A | D | S | V | A | L | H | D | E | A | A | Y | L | R | S | G | K | N | P | Y | L | F | S | P | A | D | Q | --- | 102 | (CAA77024) |
| AAV6 PV | 45 | FVLP | Y | GP | GN | S | D | --- | K | P | V | N | K | A | D | S | V | A | L | H | D | E | A | A | Y | L | R | S | G | K | N | P | Y | L | F | S | P | A | D | Q | --- | 103 | (AAB95450) |
| Bovine PV | 13 | LTLP | Y | GP | GN | S | D | --- | K | P | V | N | K | A | D | S | V | A | L | H | D | E | A | A | Y | L | R | S | G | K | N | P | Y | L | F | S | P | A | D | Q | --- | 71 | (VCPVB5) |
| Simian PV | 158 | LTLP | Y | GP | GN | S | D | --- | K | P | V | N | K | A | D | S | V | A | L | H | D | E | A | A | Y | L | R | S | G | K | N | P | Y | L | F | S | P | A | D | Q | --- | 216 | (AAA74974) |
| Rh/Macaq. PV | 125 | LTLP | Y | GP | GN | S | D | --- | K | P | V | N | K | A | D | S | V | A | L | H | D | E | A | A | Y | L | R | S | G | K | N | P | Y | L | F | S | P | A | D | Q | --- | 183 | (AAF61214) |
| Chipmunk PV | 166 | LTLP | Y | GP | GN | S | D | --- | K | P | V | N | K | A | D | S | V | A | L | H | D | E | A | A | Y | L | R | S | G | K | N | P | Y | L | F | S | P | A | D | Q | --- | 224 | (AAB82734) |
| B19 PV | 123 | LTLP | Y | GP | GN | S | D | --- | K | P | V | N | K | A | D | S | V | A | L | H | D | E | A | A | Y | L | R | S | G | K | N | P | Y | L | F | S | P | A | D | Q | --- | 181 | (VCPV19) |
| PLA2 IA | 47 | DFADY | Y | GP | GN | S | D | --- | T | V | D | L | D | R | C | Q | V | H | D | N | G | N | E | A | (X33) | AV | C | D | R | L | A | A | I | C | (X33) | 126 | (I51017) | | | | | | |
| PLA2 IB | 43 | EYNN | Y | GP | GN | S | D | --- | T | V | D | L | D | R | C | Q | V | H | D | N | G | N | E | A | (X38) | FI | C | N | C | D | R | N | A | A | I | C | (X38) | 127 | (NP_000919) | | | | |
| PLA2 IIA | 40 | SYGF | Y | GP | GN | S | D | --- | S | P | K | A | T | D | R | C | V | H | D | C | G | K | R | L | (X31) | Q | L | C | E | C | D | K | A | A | A | I | C | (X31) | 117 | (PSHUYF) | | | |
| PLA2 IIB | 18 | DYIY | Y | GP | GN | S | D | --- | K | P | V | N | K | A | D | S | V | A | L | H | D | E | A | A | Y | L | R | S | G | K | N | P | Y | L | F | S | P | A | D | Q | --- | 92 | (PSBGA) |
| PLA2 IIC | 48 | SYYG | Y | GP | GN | S | D | --- | I | P | B | A | T | D | R | C | V | H | D | C | G | K | R | L | (X28) | E | L | C | E | C | D | R | V | A | A | I | C | (X28) | 129 | (B54762) | | | |
| PLA2 III | 29 | IITY | Y | GP | GN | S | D | --- | I | P | B | A | T | D | R | C | V | H | D | C | G | K | R | L | (X35) | K | A | C | E | C | D | K | L | S | V | Y | C | (X35) | 98 | (P00630) | | | |
| PLA2 V | 40 | NKGF | Y | GP | GN | S | D | --- | T | V | D | L | D | R | C | Q | V | H | D | N | G | N | E | A | (X17) | L | S | C | E | C | D | K | F | V | D | C | (X17) | 98 | (P00630) | | | | |
| PLA2 X | 61 | AMK | Y | GP | GN | S | D | --- | T | V | D | L | D | R | C | Q | V | H | D | N | G | N | E | A | (X31) | L | L | C | K | C | D | E | I | A | N | C | (X31) | 139 | (NP_003552) | | | | |

Figure 2





A. Activity of expressed VP1up PLA₂B. Size of PPV VP1up and PLA₂ activityC. Impact Ca²⁺, pH and substrate on pvPLA₂D. PLA₂ activity of PPV virions

Figure 3




Figure 5




Figure 6

BEST AVAILABLE COPY





Figure 7

Porcine Parvovirus V1up

```

1 GCGCCTCCTGCAAAAAGAGCAAGAGGACTAACTCTACCAGGATACAAATA
51 CCTTGGTCCAGGAACTCACTAGACCAAGGAGAACCAACTAATCCATCAG
101 ACGCCGCAGCAAAAGAACACGACGAAGCCTACGACAAATACATAAAATCT
151 GGAAAAAATCCATACTTCTACTTCTCAGCAGCTGATGAAAAATTCATAAA
201 AGAAACTGAACACGCAAAAGACTACGGAGGTAAAATTGGACATTACTTCT
251 TCAGAGCAAAGCGTGCCTTTGCTCCAAAACCTCTCAGAAACAGACTCACCA
301 ACTACATCTCAACAACCAGAGGTAAAGAAGATCGCCGAGAAAACACCCAGG
351 GTCTAAACCACCAGGAAAAAGACCTGCTCCAAGACATATTTTATAAACT
401 TAGCTAAAAAAAAGCTAAAGGGACATCTAATACAACTCTAACTCAATG
451 AGTGAAAATGTGGAACAACACAACCCTATTAATGCAGGCACTGAATTGTC
501 TGCAACAGGAAATGAA

```

B19 Parvovirus V1up

```

1 AGTAAAGAAAGTGGCAAATGGTGGGAAAGTGATGATAAATTTGCTAAAGC
51 TGTGTATCAGCAATTTGTGGAATTTATAAAAAGGTTACTGGAACAGACT
101 TAGAGCTTATTCAAATATTTAAAGATCATTATAATATTTCTTTAGATAAT
151 CCCCTAGAAAACCCATCCTCTCTGTTTGACTTAGTTGCTCGTATTAAAAA
201 TAACCTTAAAACTCTCCAGACTTATATAGTCATCTTTCAAAGTCATG
251 GACAGTTATCTGACCACCCCATGCCTTATCATCCAGTAGCAGTCATGCA
301 GAACCTAGAGGAGAAAATGCAGTATTATCTAGTGAAGACT ACACAAGCC
351 TGGGCAAGTTAGCGTACAACCTACCCGGTACTAACTATGTTGGGCCTGGCA
401 ATGAGCTACAAGCTGGGCCCCCGCAAAGTGCTGTTGACAGTGCTGCAAGG
451 ATTCATGACTTTAGGTATAGCCAACCTGGCTAAGTTGGGAATAAATCCATA
501 TACTCATTGGACTGTAGCAGATGAAGAGCTTTTAAAAAATATAAAAAATG
551 AAAGTGGGTTTCAAGCACAAAGTAGTAAAGACTACTTTACTTTAAAGGT
601 GCAGCTGCCCCTGTGGCCCATTTTCAAGGAAGTTTGCCGGAAGTTCCCGC
651 TTACAACGCCTCAGAAAAATACCCAAGC

```

Galleria mellonella Densovirus V1up

```

1 ATGTCCTTCTTCAAAAATCAGTTGATACATCGCGCACGACCTGGTTATCG
51 TATAATACCGGAAAGTACTGTTACTGAAGATATTGAATTAGGTACTATTG
101 GTGAAGAAACTCCATTGTTAAGTGAAGGTGTTATTACAGCTGTAGAAGAA
151 GGTGCTATTGGATTACCAGAAGTTGCTATTGGTGTGGCTGGTGTCTATTGG
201 AACACATGCACATGAATGGTGGAGAGATAGATACGCTTTTAAATCTGTTT
251 TAACTGGTAATTATACTGATTTTAAAGGAAATCCTTTAAACCTAGAAAC
301 GCTATTCCTGAAAAAATTAACAACCTAGGAAAGAAAATATTTCAAGGAGA
351 TTTTAATCGTGCATTTCTGATAATTTAAAATTGGAAACTGAAAAAGAAA
401 AAGCTGATTTATTAAGATATTATAATCATAATAGAAGATTAGCTGGACTA
451 AGTGAAGCTTATCCACAAGGGAAAGGATATGCTTATGCTAAAAGTCAAAA
501 AGTATTAGAAGCTGAACGACGTGGATTAAGTGTTCCTGGATATAAATATC
551 TTGGTCCTGGAAATTCATTGAATAGAGGTCAACCTATAAATCAAATAGAC
601 GAAGACGCTAAAGAACACGACGAAGCTTATGATAAAGTGAAAAACAAGTCA
651 AGAAGTAAAGTCGAGCAGATAATACATTTGTTAATAAAGCGTTAGATCACG
701 TGGTTAATGCTATTAATTTTAAAGAAACACCTGGTAACGCTTTTGGTGCT
751 GCTATTGGAGCTATTGGAATTGGAACCTAAGCAAGCTATTGAAAAATATAG
801 TGGAGTAATCTACCTTCTGTTTCAGGT

```

Figure 8

***Bombyx mori* Densovirus V1up**

1 ATGCCTCGTATTCATTTTCCTTATCATAATTATCTTGGTCCGGGTACTGA
51 TAACTTTGAATAAAATCCAATAGACGAAGACGACGCTATCGCGAGGTCCG
101 ATGATTTGGCTTACGATAAAGTAACCAATCATAAGGAAGTTTTTCAAGCT
151 GATAAACAGGCCCGTGACGAGTTTTTTACTTCATTTGTGCATACTGGAAA
201 CGTGCATAGTTTAATTGGCGGTATTGGACTTGGAACTAAAAATTTGGTAG
251 AAGAACATGTACTAGGTAAACCTTGTACGGA

***Casphalia extranea* Densovirus V1up**

1 ATGCCTCGTATTCATTTTCCTTATCATAATTATCTTGGTCCGGGCAGTGA
51 TAACTTTAAAAACAACCAGTAGACGAAGACGACGCAATAGCCAGAGCAC
101 ATGACCTGGATTACGATAAAGCAAGCTCTGATAAAGACATTTTCAAGGCT
151 GATAAGCAGGCTCGCGACGAGTTTTTCAGTTTCATTTGTGCACAGCGGAAA
201 CTTGCATAGTTTAATTGGTGGACTAGGACTTGGAGCTAAAAATCTAGTAG
251 AAGAGCATGTACTAGGTAAAGTCCTTGTACGGT

Adeno-Associated Virus V1up

1 ATGGCTGCCGATGGTTATCTTCCAGATTGGCTCGAGGACACTCTCTCTGA
51 AGGAATAAGACAGTGGTGGAAAGCTCAAACCTGGCCCACCACCACCAAAGC
101 CCGCAGAGCGGCATAAGGACGACAGCAGGGGTCTTGTGCTTCTGGGTAC
151 AAGTACCTCGGACCCTTCAACGGA CTGACAAGGGAGAGCCGGTCAACGA
201 GGCAGACGCCGCGGCCCTCGAGCACGACAAAGCCTACGACCGGCAGCTCG
251 ACAGCGGAGACAACCCGTACCTCAAGTACAACCACGCCGACGCGGAGTTT
301 CAGGAGCGCCTTAAAGAAGATACGTCCTTTTGGGGGCAACCTCGGACGAGC
351 AGTCTTCCAGGCGAAAAAGAGGGTTCTTGAACCTCTGGGCCTGGTTGAGG
401 AACCTGTTAAGACGGCTCCGGGAAAAAAGAGGCCGGTAGAGCACTCTCCT
451 GTGGAGCCAGACTCCTCCTCGGGAACCGGAAAGGCCGGGCCAGCAGCCTGC
501 AAGAAAAAGATTGAATTTTGGTCAGACTGGAGACGCAGACTCAGTACCTG
551 ACCCCCAGCCTCTCGGACAGCCACCAGCAGCCCCCTCTGGTCTGGGAACT
601 AATACG

Figure 8 (con.)

Porcine Parvovirus V1up

1 APPAKRARGLTLPGYKYLGPNSLDQGEPTNPSDAAAKEHDEAYDKYIKS
 51 GKNPYFYFSAADEKFIKETEHAADYGGKIGHYFFRAKRAFAPKLSETDSP
 101 TTSQQPEVRRSPRKHPGSKPPGKRPAPRHIFINLAKKKAKGTSNTNSNSM
 151 SENVEQHNPINAGTELSATGNE

B19 Parvovirus V1up

1 SSKEGKWWESDDKFAKAVYQQFVEFYKKVTGTDLELIQILKDHYNISLD
 51 NPLENPSSFLDLVARIKNNLNKNSPDLYSHHFQSHGQLSDHPPHALSSSSSH
 101 AEPRGENAVLSSDLHKPGQVSVQLPGTNYVGPNGELQAGPPQ\$AVDSAA
 151 RIHDFRYSQLAKLGINPYTHWTVADEELLKNIKNETGFQAQVVKDYFTLK
 201 GAAAPVAHFQGSLEVPAYNASEKYPSAYNASEKYPS

Galleria mellonella Densovirus V1up

1 MSFFKNQLIHRARPGYRIIPESTVTEDIELGTIGEETPLLSEGVITAVEE
 51 GAIGLPEVAIGVAGAIGTHAHEWWRDRYAFKSVLTGNYTDLKGNPLKPRN
 101 AIPEKIKQLGKKIFQGDFNRAFPDNLKLETEKEKADLLRYNHNRRLAGL
 151 SEAYPQGKGYAYAKSQKVLEAERRGLTVPGYKYLGPNSLNRGQPINQID
 201 EDAKEHDEAYDKVKTSQEVSRADNTFVNKALDHVVNAINFKETPGNAFGA
 251 AIGAIGIGTKQAIKYSYSGVIYPSVSG

Bombyx mori Densovirus V1up

1 MPRIHFYPYHNYLGPSTDNFE KNPIDEDDAIARSHDLAYDKVTNHKEVFQA
 51 DKQARDEFFTSFVHTGNVHSLIGGIGLGTKNLVEEHVLGKPLYG

Casphalia extranea Densovirus V1up

1 MPRIHFYPYHNYLGPSTDNFKNQPVDEDDAIARAHDLDYDKASSDKDIFKA
 51 DKQARDEFFSSFVHSGNLHSLIGGLGLGAKNLVEEHVLGKSLYG

Adeno-associated Virus V1up

1 MAADGYLPDWLEDLSEGIQWWKLKPGPPPPKPAERHKDDSRGLVLPGY
 51 KYLGPFNGLDKGEPVNEADAAAEHDKAYDRQLDSGDNPYLKYNHADAEF
 101 QERLKEDTSFGGNLGRAVFQAKKRVLEPLGLVEEPVKTA PGKKRPVEHSP
 151 VEPDSSSGTGKAGQQPARKRLNFGQTGDADSVDPQPLGQPPAAPSGLGT
 201 NT

Figure 9

| | | | | | |
|------|-------------|-------------|------------|-------------|-------------|
| 1 | TTATATCCTT | AGGATATRAA | GAGCAAGCAC | CCAGGCACCTT | CTTCTATTCT |
| 51 | GCCTCAGATC | GTCCGATATRA | CGCGAGCTTT | TCTGTGACCTG | TAGGTAGTGT |
| 101 | ATTGATCTTC | GTGTTCTGGA | ACTCAGGTAT | GTGTTTCGCT | GCAGTAGAAG |
| 151 | CGCTCTCAGC | ATGACGTGTC | GAACCTCGTT | CATTGCAACA | GTATGACATG |
| 201 | AGAACTTTTG | TAGAAATATC | GGATTGGGAC | GTATATTAAT | TTCTCTATAA |
| 251 | AAATATCTGA | TTTCTTAGCA | GSTTGTATTC | ATCGATCTTT | TATGARAAGA |
| 301 | AAATATCTGA | TAATCTCGCAT | GAATATCATG | ATTTGGTCAGT | ACCAATATAT |
| 351 | TTTCCCATTC | AAACGTATAT | AAAGATATAT | AAATATCACA | TGTTGATACG |
| 401 | TATTCATATC | TTTGTCTTCC | RACGCAATTT | AGTGAATAGA | TGTTTATCAT |
| 451 | AAATGCAATTT | CTGTGCTCCT | GSTACTTTTG | GCTTCCCTAT | TACAGATATAT |
| 501 | ATTGATATAC | TCTGTATATA | TACATTTTGG | AGGTTTAAAC | GCCTTATATTT |
| 551 | CTATCGGTTT | AAAGATTTTG | TTCTCTGTCG | CGATCAACAA | GTATTCTAAG |
| 601 | ATGATGAACT | TATCGATTTAT | ATGCAATGTA | GAATTAATTG | GTCTACTTGT |
| 651 | TGTAATACAA | CATCATTTAT | TCCGATTAAT | GAATGAACTC | ATTAGGCCCC |
| 701 | GCATGCTGTT | AATGATATAT | ATCTCATGTC | TACGCAATTT | CCGTTATTAT |
| 751 | CTGTTGCAAG | TGATRACTGA | GTAAACGAT | GGAGCCAGTA | CRAAGCACAGG |
| 801 | ACATCTGAGA | TGGAGTAGAG | GAGACCTTGA | AGACCTTTAT | GAAGACATTA |
| 851 | TGGGCCCTCT | CGAGAATCA | ACCGTTGACC | AAATCCGCAAT | GGTGGGCGGA |
| 901 | CATGCTGAG | AGTGTATCCA | TTTCTCTGGA | GGATCAGAAC | ACCAATCTAG |
| 951 | CCTCAGATTA | CGCAAAATTA | AGAAAGCT | TCTGGAAGT | TATATCCAC |
| 1001 | TGGCAGACAA | TGTCACGAA | ATGTTTGGTC | ACTTCAATCG | CGACCTTAGA |
| 1051 | TCGAAGATTT | TCRGCTCTAC | TGGGGGTAC | ATTTCAGCTC | TTGTGCTCAC |
| 1101 | GTGCTCTCCT | GAACACGGTC | CGAGGTGCT | GGACTTCTTT | ATAAGTGGC |
| 1151 | CAACGACTTA | CCCCGGAAGA | CTACTACTCT | GGTCCACAG | AGGACACCA |
| 1201 | ATTCACGCTG | TCCAGACATG | CCCTACAGC | GCTGGACAA | GTAGATGCAA |
| 1251 | GTCTCTAA | ACGGAAGCT | TCAAAAGGG | ACTTCGCAAG | CCCTACGAA |
| 1301 | ATGCAAGTT | CATTACCGAA | CTGCTAGCA | CGACCTGGGC | AAATGTTTTC |
| 1351 | TTTACTTTCA | TTGCTCAAA | ATGGBAAAC | GAACGAGCAG | TTTGGATTGA |
| 1401 | TGGAGCCTTA | CGACGACTAC | CGGATACAG | TGAAGGTGTA | CGATGGACCG |
| 1451 | CGCTGTGGA | GCAATCCCGA | CTACTACTGG | AACACAGAGC | TCGAACAGAT |
| 1501 | GGATGTGAG | GTATCGAAGA | ATCTACAGAT | TTGGAAGGAC | ATCAACAGAG |
| 1551 | CGTTTGTGCT | GGCGTATACC | AACTCCGAG | AAACGGAGC | AGTTCCAGA |
| 1601 | CTTTCGCGGA | AGAAGCAAAA | GTACTATTCT | CAAAATTTGA | CTGTATTCCT |
| 1651 | GTGCTGAGC | CTAGAAATAT | AAATGATAGG | AAACATCTTA | TGTATTTTGA |
| 1701 | ATTGCATGAT | CCAAAATATG | AAAAGTTGTA | TACTGTGCT | ATTCTTTAT |
| 1751 | ATGTCGCTGA | CATTATGAA | ATGCTTTGA | ATGATTTCTA | CTTACAGACT |
| 1801 | TTAGAGAAAT | CATGTGTTTA | TCAAGCTGTT | AAATATAGCC | CTTGTGTTT |
| 1851 | TTATCATATAT | ATGGAAGATA | GTGTTACAT | TCTTATGAA | CTTATCTTAT |
| 1901 | TTCAATGTAA | TGGTAGTAG | GAAGTGTGA | AAATATCT | ATGAATATTT |
| 1951 | TGAGCTTGGT | TAATATGTT | AGGTTGGTAT | ATGCCAACAA | TTAATATGAA |
| 2001 | TGGGAAATA | GAATATCAG | TAAACCTTAA | AGTTATAGCT | GTATGTATTA |
| 2051 | TTGGAATCA | TAATGTGTT | AGATTTAT | TTTGGACGC | AGTTGCTGT |
| 2101 | TTGGTTTAA | ATGTTGGTT | TCTTGTGTA | GTATTAATA | AAATATACAA |
| 2151 | TTTGTCTTGG | CAAGATTTG | TTCAAAAGC | AAATGTAAT | GTATATGAAA |
| 2201 | TATCTTTTGA | AGATGAGCA | RAGGAGATA | TGAAGAAAT | ATGTGAGGG |
| 2251 | TGCTCATTTCA | ATGTTAGCT | AAACATCAA | TCAGATGAA | TGTTGCTCAG |
| 2301 | GACACCTTTT | TGCTTAATA | GTACAAATG | TTTACTTAT | GAATGATG |
| 2351 | CTCATTTTGT | AGATTAGAG | ATGAGGAT | TTTATTTAG | CGATGATG |
| 2401 | TATTTTGGCA | ATGCAAGAA | GAAGCGGTAT | CCACCGGCT | TTTCTTTAT |
| 2451 | AATGAAATG | TATGAGTTG | ATTGGAATA | AGTTTATTA | TTTCTTTAT |
| 2501 | TACATCTTTC | ATTTTGTG | CAATCCCAT | ATAGTACTCT | TCCCACTGC |
| 2551 | ATGATGTGAC | TCCAAAGTAG | AACTGGAATA | AACTGCTTA | TCAGGAATAG |
| 2601 | TATTAGGACC | GTAAANAGA | GCTCTAATGG | AAAGATATTG | AAATGCTACT |
| 2651 | TTGCRAGCAG | CTTCAACAAT | AAATGATAGT | TGAGTGTGAC | TAAATATTAGC |
| 2701 | ATATATTAGA | TATATCACTA | ANTATGCTGT | TGTCANATGA | TGACACAGTT |
| 2751 | TACACCAAT | ATGCAACTA | GGTGTAGCTT | TTGTTGTATA | AGATGGAATA |
| 2801 | GTCTCTCAGC | GAATATTTG | ACTTTTCTCA | ATTTGAGTCC | CAATATGCTCA |
| 2851 | TTTCAATTTT | GATATATCAA | TATCATTTAA | TTTTTCTGAA | TCCCTCTTCA |
| 2901 | AAATATCTTT | AGTCAATAGT | AAATCAAGC | TCACCTACTG | CGATGCTCCT |
| 2951 | AAATATCTTT | AGTCAATAGT | AAATCAAGC | TCACCTACTG | CGATGCTCCT |
| 3001 | AAATATCTTT | AGTCAATAGT | AAATCAAGC | TCACCTACTG | CGATGCTCCT |
| 3051 | AAATATCTTT | AGTCAATAGT | AAATCAAGC | TCACCTACTG | CGATGCTCCT |
| 3101 | AAATATCTTT | AGTCAATAGT | AAATCAAGC | TCACCTACTG | CGATGCTCCT |
| 3151 | AAATATCTTT | AGTCAATAGT | AAATCAAGC | TCACCTACTG | CGATGCTCCT |
| 3201 | AAATATCTTT | AGTCAATAGT | AAATCAAGC | TCACCTACTG | CGATGCTCCT |
| 3251 | AAATATCTTT | AGTCAATAGT | AAATCAAGC | TCACCTACTG | CGATGCTCCT |
| 3301 | AAATATCTTT | AGTCAATAGT | AAATCAAGC | TCACCTACTG | CGATGCTCCT |
| 3351 | AAATATCTTT | AGTCAATAGT | AAATCAAGC | TCACCTACTG | CGATGCTCCT |
| 3401 | AAATATCTTT | AGTCAATAGT | AAATCAAGC | TCACCTACTG | CGATGCTCCT |
| 3451 | AAATATCTTT | AGTCAATAGT | AAATCAAGC | TCACCTACTG | CGATGCTCCT |
| 3501 | AAATATCTTT | AGTCAATAGT | AAATCAAGC | TCACCTACTG | CGATGCTCCT |
| 3551 | AAATATCTTT | AGTCAATAGT | AAATCAAGC | TCACCTACTG | CGATGCTCCT |
| 3601 | AAATATCTTT | AGTCAATAGT | AAATCAAGC | TCACCTACTG | CGATGCTCCT |
| 3651 | AAATATCTTT | AGTCAATAGT | AAATCAAGC | TCACCTACTG | CGATGCTCCT |
| 3701 | AAATATCTTT | AGTCAATAGT | AAATCAAGC | TCACCTACTG | CGATGCTCCT |
| 3751 | AAATATCTTT | AGTCAATAGT | AAATCAAGC | TCACCTACTG | CGATGCTCCT |
| 3801 | AAATATCTTT | AGTCAATAGT | AAATCAAGC | TCACCTACTG | CGATGCTCCT |
| 3851 | AAATATCTTT | AGTCAATAGT | AAATCAAGC | TCACCTACTG | CGATGCTCCT |
| 3901 | AAATATCTTT | AGTCAATAGT | AAATCAAGC | TCACCTACTG | CGATGCTCCT |
| 3951 | AAATATCTTT | AGTCAATAGT | AAATCAAGC | TCACCTACTG | CGATGCTCCT |
| 4001 | AAATATCTTT | AGTCAATAGT | AAATCAAGC | TCACCTACTG | CGATGCTCCT |
| 4051 | AAATATCTTT | AGTCAATAGT | AAATCAAGC | TCACCTACTG | CGATGCTCCT |
| 4101 | AAATATCTTT | AGTCAATAGT | AAATCAAGC | TCACCTACTG | CGATGCTCCT |
| 4151 | AAATATCTTT | AGTCAATAGT | AAATCAAGC | TCACCTACTG | CGATGCTCCT |
| 4201 | AAATATCTTT | AGTCAATAGT | AAATCAAGC | TCACCTACTG | CGATGCTCCT |
| 4251 | AAATATCTTT | AGTCAATAGT | AAATCAAGC | TCACCTACTG | CGATGCTCCT |
| 4301 | AAATATCTTT | AGTCAATAGT | AAATCAAGC | TCACCTACTG | CGATGCTCCT |
| 4351 | AAATATCTTT | AGTCAATAGT | AAATCAAGC | TCACCTACTG | CGATGCTCCT |
| 4401 | AAATATCTTT | AGTCAATAGT | AAATCAAGC | TCACCTACTG | CGATGCTCCT |
| 4451 | AAATATCTTT | AGTCAATAGT | AAATCAAGC | TCACCTACTG | CGATGCTCCT |
| 4501 | AAATATCTTT | AGTCAATAGT | AAATCAAGC | TCACCTACTG | CGATGCTCCT |
| 4551 | AAATATCTTT | AGTCAATAGT | AAATCAAGC | TCACCTACTG | CGATGCTCCT |
| 4601 | AAATATCTTT | AGTCAATAGT | AAATCAAGC | TCACCTACTG | CGATGCTCCT |
| 4651 | AAATATCTTT | AGTCAATAGT | AAATCAAGC | TCACCTACTG | CGATGCTCCT |
| 4701 | AAATATCTTT | AGTCAATAGT | AAATCAAGC | TCACCTACTG | CGATGCTCCT |
| 4751 | AAATATCTTT | AGTCAATAGT | AAATCAAGC | TCACCTACTG | CGATGCTCCT |
| 4801 | AAATATCTTT | AGTCAATAGT | AAATCAAGC | TCACCTACTG | CGATGCTCCT |
| 4851 | AAATATCTTT | AGTCAATAGT | AAATCAAGC | TCACCTACTG | CGATGCTCCT |
| 4901 | AAATATCTTT | AGTCAATAGT | AAATCAAGC | TCACCTACTG | CGATGCTCCT |
| 4951 | AAATATCTTT | AGTCAATAGT | AAATCAAGC | TCACCTACTG | CGATGCTCCT |
| 5001 | AAATATCTTT | AGTCAATAGT | AAATCAAGC | TCACCTACTG | CGATGCTCCT |
| 5051 | AAATATCTTT | AGTCAATAGT | AAATCAAGC | TCACCTACTG | CGATGCTCCT |
| 5101 | AAATATCTTT | AGTCAATAGT | AAATCAAGC | TCACCTACTG | CGATGCTCCT |
| 5151 | AAATATCTTT | AGTCAATAGT | AAATCAAGC | TCACCTACTG | CGATGCTCCT |
| 5201 | AAATATCTTT | AGTCAATAGT | AAATCAAGC | TCACCTACTG | CGATGCTCCT |

Figure 10

***Acheta domesticus* Densovirus nucleic acid sequence containing PLA₂ motif**

```

1  ATGTCTGGCG TCTTTACAGA TCTCACGTTG CGTGATTTCA GTTTAGGTAC
51 TAATAATTTA CCTTTAGCAA ATACACCTAA ACTTCGTAAT AGATTTGGGA
101 GATGGTGGAA TCGTAGTCAC CCCTACGACA GACTACCTAC AAATGAACCA
151 GAACCTCTTC GCGAAACTAG TTTCCAAGAG GCGGCAGGGC CAGAAGAAAC
201 TCGCATTGAT ATCGCGGAAG ACGAAATCAA CGCAGGTGAA GGAGCAGCCG
251 AAGCTGAAAC AAGTTTCTCT ACTGGAGTTG AGGAAACAGC ACTTGAACTC
301 GGCGAAGCAG CTACCGAAAC AACAGGACTT TTGGGAGGCG CTACTGCAGG
351 TAGCGCAGCA GCAGGCACAG CTGGAACCTCT TGGAACTGTC GCGGCAACTG
401 CAGCAGGAGG AGCAGCCCTC GCAGGAATCG GCATCGGAAT TAAAAAATTA
451 ATTGATCACA CGAGTAGCAA AGGTGCAGTA TTACCTGGCA CTGATTTCTGT
501 GGGGCCTGGT AATCCAATAG ATCCTAAACC TGCACGATCA GAAACTGATC
551 AAATTGCTAA GGAACACGAT CTTGGGTACG AAGACTTATT GCATCGTGCA
601 AAATCACAGT ATTTTACTGA AGAAGATTTT AAAACTGAAG TATATAAATT
651 GGACGACGAA GCAATTCATC GTTTTCTGA GGAATACCAG AAATCCGGTA
701 CGTGCCAAGC ATTTGTTGGA AAATACGGAC TTAAAGCTAA ACGTGTAATT
751 GAAGACGTAA TTGGTGGACC TGTATATCCT CAACAACCTA AAAACGTGA
801 GTAC

```

***Mythimna loreyi* Densovirus nucleic acid sequence containing PLA₂ motif**

```

1  ATGTCTTTCT ACACGAGCGG GTTAACACAT CGCGCGCGAC CTGGCTATCG
51 TATAATACCA GAAAGTACTG CTACCGAAGA TATAGAATTA GGTACTATTG
101 GTGAAGAAAC TCCTTTATTA AGTGAAAGTG CTATTACTGC TGTAGAAGAA
151 GGTGCTATTG GAGTTCCAGA AGTTGCCGTT GGAATTGCTG GTGCTATAGG
201 AACACATGCT GACGTGTTAT ATAGAAATAG AAACGTATTT AAAAGTGTTT
251 TAACTGGAAA TTACACTGAT TTAAAAGGAA ATCCAATAAA ACAAAGAAAT
301 TCTATTCCTG AGAAGACAAA GCAATTAGGA AAAGGTATAT TTCAAGGTGA
351 TTTTAATCGT GCATTCCTG AAGATTTAAA AACAGAACT GAGCAAGAAA
401 AAAACGATTT ATTACGATAC TATAATCATA ATAGAAGACT AGCTGGTTTA
451 AGTGAAGCTT ATCCACAAGG AAAAGGATAC GCTTATGCTA AGAGTCAAAA
501 AGTATTAGAA GCTGAAAGAC GTGGATTAAAC TGTTCTGGA TATAAATATC
551 TTGGTCCTGG AAACCACTT AATAGAGGTC AACCTACTAA TCAAATAGAC
601 GAAGACGCTA AAGAACACGA CGAAGCTTAC GATAAAGCAA AAACAAGTCA
651 AGAAGTAAGT GAAGCAGATA ATACATTTGT TAATAAGGCG TTAGATCACG
701 TGGTTAATGC TATCAATTTT AAAGAAACGC CTGGTAACGC TTTTGGTGCT
751 GCTATTGGAG CTATTGGAAT TGGAACTAAG CAAGCTATTG AAAACACAG
801 TGGAGTAATC TACCCTTCTG TTTCAGGT

```

Figure 11

***Acheta domesticus* Densovirus amino acid sequence containing PLA₂ motif**

1 MSGVFTDLTL RDFS LGTNNL PLANTPKLRN RFGRWWNRSH PYDRLPTNEP
51 EPLRETSFQE AAGPEETRID IAED EINAGE GAAEAETSFS TGVEETALEL
101 GEAATETTGL LGGATAGSAA AGTAGTLGTV AATAAGGAAL AGIGIGIKKL
151 IDHTSSKGAV LPGTDFVGPG NPIDPKPARS ETDQIAKEHD LGYEDLLHRA
201 KSQYFTEEDF KTEVYKLDDE AIHRFSE EYQ KSGTWQAFVG KYGLKAKRVI
251 EDVIGGPVYP QQPKKREY

***Mythimna loreyi* Densovirus amino acid sequence containing PLA₂ motif**

1 MSFYTSGLTH RARPGYRIIP ESTATEDIEL GTIGEETPLL SESAITAVEE
51 GAIGVPEVAV GLAGAIGTHA DVLYRNRNVF KSVLTGNYTD LKGNPIKQRN
101 SIPEKTKQLG KGIFQGD FNR AFPEDLKTET EQEKNDLLRY YNHNRRLAGL
151 SEAYPOGKGY AYAKSQK VLE AERRGLTVPG YKYLGP GNSL NRGQPTNQID
201 EDAKEHDEAY DKAKTSQEV S EADNTFVNKA LDHVVNAINF KETPGNAFGA
251 AIGAIGIGTK QAIEKHSGVI YPSVSG

Figure 12

P7

DNA: TGGTCTAGTTCGCATTATCCTCCGCATTGGACTGCG
+1: W S S S H Y P P H W T A

P6

DNA: GTGAATCAGTCTTATACTTCGACTTGGTTTGGGCT
+1: V N Q S Y T S T W F W A

P8

DNA: TGGCCTCAGTTGTTTACTTTTCCGTGTTGTAATCCT
+1: W P Q L F T F P C C N P

P10

DNA: GGTCTGAAGATTGGAGTTTGCCGCCGCATCATGGG
+1: G L K I W S L P P H H G

Figure 13

14/14

SEQUENCE LISTING

<110> Tjissen, Peter
Zadori, Zoltan

<120> VIRAL PHOSPHOLIPASE A2 ENZYMES,
ANTI-VIRAL AGENTS AND METHODS OF USE

<130> 255-123pct

<140> unavailable

<141> 2001-06-28

<150> 2,310,263

<151> 2000-06-28

<160> 73

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 54

<212> PRT

<213> Galleria mellonella densovirus

<400> 1

```

Leu Thr Val Pro Gly Tyr Lys Tyr Leu Gly Pro Gly Asn Ser Leu Asn
 1             5             10             15
Arg Gly Gln Pro Ile Asn Gln Ile Asp Glu Asp Ala Lys Glu His Asp
      20             25             30
Glu Ala Tyr Asp Lys Val Lys Thr Ser Gln Glu Val Ser Arg Ala Asp
      35             40             45
Asn Thr Phe Val Asn Lys
      50

```

<210> 2

<211> 54

<212> PRT

<213> Mythimna loreyi densovirus

<400> 2

```

Leu Thr Val Pro Gly Tyr Lys Tyr Leu Gly Pro Gly Asn Ser Leu Asn
 1             5             10             15
Arg Gly Gln Pro Thr Asn Gln Ile Asp Glu Asp Ala Lys Glu His Asp
      20             25             30
Glu Ala Tyr Asp Lys Ala Lys Thr Ser Gln Glu Val Ser Glu Ala Asp
      35             40             45
Asn Thr Phe Val Asn Lys
      50

```

<210> 3
 <211> 54
 <212> PRT
 <213> Junonia coenia densovirus

<400> 3
 Leu Thr Val Pro Gly Tyr Lys Tyr Leu Gly Pro Gly Asn Ser Leu Asn
 1 5 10 15
 Arg Gly Gln Pro Thr Asn Gln Ile Asp Glu Asp Ala Lys Glu His Asp
 20 25 30
 Glu Ala Tyr Asp Lys Ala Lys Thr Ser Gln Glu Val Ser Gln Ala Asp
 35 40 45
 Asn Thr Phe Val Asn Lys
 50

<210> 4
 <211> 54
 <212> PRT
 <213> Pseudoplusia includens densovirus

<400> 4
 Leu Thr Val Pro Gly Tyr Lys Tyr Leu Gly Pro Gly Asn Ser Leu Asp
 1 5 10 15
 Arg Gly Glu Pro Val Asn Gln Ile Asp Ala Asp Ala Lys Glu His Asp
 20 25 30
 Glu Ala Tyr Asp Lys Ala Lys Thr Ser Gln Glu Val Ser Asp Ala Asp
 35 40 45
 Ser Lys Phe Val Ser Lys
 50

<210> 5
 <211> 54
 <212> PRT
 <213> Diatraea saccharalis densovirus

<400> 5
 Leu Thr Val Pro Gly Tyr Lys Tyr Leu Gly Pro Gly Asn Ser Leu Asn
 1 5 10 15
 Arg Gly Pro Pro Thr Asn Glu Ile Asp Ala Asp Ala Lys Glu His Asp
 20 25 30
 Glu Ala Tyr Ser Gln Ser Lys Thr Ala Gln Glu Val Ser Lys Ala Asp
 35 40 45
 Asn Thr Phe Val Asn Lys
 50

<210> 6
 <211> 54
 <212> PRT
 <213> Culex pipiens densovirus

<400> 6

Leu Val Pro Ala Pro Tyr Lys Tyr Ala Gly Pro Gly Asn Ser Leu Asn
 1 5 10 15
 Arg Gly Pro Ala Tyr Asp Leu Val Asp Glu Ser Ala Arg Gln His Asp
 20 25 30
 Ile Ala Tyr Asp Lys Ala Lys Ser Pro Glu Asp Ile His Lys Ala Asp
 35 40 45
 Arg Gln Phe Leu Thr Glu
 50

<210> 7

<211> 54

<212> PRT

<213> Periplaneta fuliginos densovirus

<400> 7

Leu Thr Tyr Pro Phe His His Tyr Leu Gly Pro Gly Asn Pro Leu Asp
 1 5 10 15
 Asn Asn Glu Pro Val Asp Arg Asp Asp Ala Ile Ala Glu Glu His Asp
 20 25 30
 Lys Ala Tyr Ala Asn Ala Lys Ser Ser Ile Asp Val Ile Asn Ala Asp
 35 40 45
 Lys Lys Ala Ile Asp His
 50

<210> 8

<211> 57

<212> PRT

<213> Acheta domesticus densovirus

<400> 8

Ala Val Leu Pro Gly Thr Asp Phe Val Gly Pro Gly Asn Pro Ile Asp
 1 5 10 15
 Pro Lys Pro Ala Arg Ser Glu Thr Asp Gln Ile Ala Lys Glu His Asp
 20 25 30
 Leu Gly Tyr Glu Asp Leu Leu His Arg Ala Lys Ser Gln Tyr Phe Thr
 35 40 45
 Glu Glu Asp Phe Lys Thr Glu Val Tyr
 50 55

<210> 9

<211> 59

<212> PRT

<213> Casphalia extranea densovirus

<400> 9

Ile His Phe Pro Tyr His Asn Tyr Leu Gly Pro Gly Ser Asp Asn Phe
 1 5 10 15
 Lys Lys Gln Pro Val Asp Glu Asp Asp Ala Ile Ala Arg Ala His Asp

```

<400> 12
Leu Val Pro Pro Gly Tyr Lys Tyr Leu Gly Pro Gly Asn Ser Leu Asp
 1          5          10          15
Gln Gly Glu Pro Thr Asn Pro Ser Asp Ala Ala Ala Lys Glu His Asp
          20          25          30
Glu Ala Tyr Ala Ala Tyr Leu Arg Ser Gly Lys Asn Pro Tyr Leu Tyr
          35          40          45
Phe Ser Pro Ala Asp Gln Arg Phe Ile Asp Gln

```

50

55

<210> 13

<211> 59

<212> PRT

<213> Mouse parvovirus 1

<400> 13

```

Leu Val Pro Pro Gly Tyr Lys Tyr Leu Gly Pro Gly Asn Ser Leu Asp
 1           5           10           15
Gln Gly Glu Pro Thr Asn Pro Ser Asp Ala Ala Ala Lys Glu His Asp
           20           25           30
Glu Ala Tyr Ala Ala Tyr Leu Arg Ser Gly Lys Asn Pro Tyr Leu Tyr
           35           40           45
Phe Ser Pro Ala Asp Gln Arg Phe Ile Asp Gln
 50           55

```

<210> 14

<211> 59

<212> PRT

<213> Feline panleukopenia parvovirus

<400> 14

```

Leu Val Pro Pro Gly Tyr Lys Tyr Leu Gly Pro Gly Asn Ser Leu Asp
 1           5           10           15
Gln Gly Glu Pro Thr Asn Pro Ser Asp Ala Ala Ala Lys Glu His Asp
           20           25           30
Glu Ala Tyr Ala Ala Tyr Leu Arg Ser Gly Lys Asn Pro Tyr Leu Tyr
           35           40           45
Phe Ser Pro Ala Asp Gln Arg Phe Ile Asp Gln
 50           55

```

<210> 15

<211> 59

<212> PRT

<213> Minute virus of mice

<400> 15

```

Met Val Pro Pro Gly Tyr Lys Tyr Leu Gly Pro Gly Asn Ser Leu Asp
 1           5           10           15
Gln Gly Glu Pro Thr Asn Pro Ser Asp Ala Ala Ala Lys Glu His Asp
           20           25           30
Glu Ala Tyr Asp Gln Tyr Ile Lys Ser Gly Lys Asn Pro Tyr Leu Tyr
           35           40           45
Phe Ser Ala Ala Asp Gln Arg Phe Ile Asp Gln
 50           55

```

<210> 16

<211> 59
 <212> PRT
 <213> Kilham rat parvovirus

<400> 16
 Cys Val Pro Pro Gly Tyr Lys Tyr Leu Gly Pro Gly Asn Ser Leu Asp
 1 5 10 15
 Gln Gly Glu Pro Thr Asn Pro Ser Asp Ala Ala Ala Lys Glu His Asp
 20 25 30
 Leu Ala Tyr Asp Glu Tyr Ile Lys Ser Gly Lys Asn Pro Tyr Leu Tyr
 35 40 45
 Phe Ser Pro Ala Asp Gln Arg Phe Ile Asp Gln
 50 55

<210> 17
 <211> 59
 <212> PRT
 <213> Porcine parvovirus

<400> 17
 Leu Thr Leu Pro Gly Tyr Lys Tyr Leu Gly Pro Gly Asn Ser Leu Asp
 1 5 10 15
 Gln Gly Glu Pro Thr Asn Pro Ser Asp Ala Ala Ala Lys Glu His Asp
 20 25 30
 Glu Ala Tyr Asp Lys Tyr Ile Lys Ser Gly Lys Asn Pro Tyr Phe Tyr
 35 40 45
 Phe Ser Ala Ala Asp Glu Lys Phe Ile Lys Glu
 50 55

<210> 18
 <211> 59
 <212> PRT
 <213> Muscovy duck parvovirus

<400> 18
 Phe Val Leu Pro Gly Tyr Lys Tyr Val Gly Pro Gly Asn Gly Leu Asp
 1 5 10 15
 Lys Gly Pro Pro Val Asn Lys Ala Asp Ser Val Ala Leu Glu His Asp
 20 25 30
 Lys Ala Tyr Asp Gln Gln Leu Lys Ala Gly Asp Asn Pro Tyr Ile Lys
 35 40 45
 Phe Lys His Ala Asp Gln Glu Phe Ile Asp Asn
 50 55

<210> 19
 <211> 59
 <212> PRT
 <213> Goose parvovirus

<400> 19

Phe Val Leu Pro Gly Tyr Lys Tyr Leu Gly Pro Gly Asn Gly Leu Asp
 1 5 10 15
 Lys Gly Pro Pro Val Asn Lys Ala Asp Ser Val Ala Leu Glu His Asp
 20 25 30
 Lys Ala Tyr Asp Gln Gln Leu Lys Ala Gly Asp Asn Pro Tyr Ile Lys
 35 40 45
 Phe Asn His Ala Asp Gln Asp Phe Ile Asp Ser
 50 55

<210> 20

<211> 59

<212> PRT

<213> Bovine parvovirus

<400> 20

Leu Thr Leu Pro Gly Tyr Asn Tyr Leu Gly Pro Phe Asn Ser Leu Phe
 1 5 10 15
 Ala Gly Ala Pro Val Asn Lys Ala Asp Ala Ala Ala Arg Lys His Asp
 20 25 30
 Phe Gly Tyr Ser Asp Leu Leu Lys Glu Gly Lys Asn Pro Tyr Leu Tyr
 35 40 45
 Phe Asn Thr His Asp Gln Asn Leu Ile Asp Glu
 50 55

<210> 21

<211> 59

<212> PRT

<213> Simian parvovirus

<400> 21

Leu Thr Leu Pro Phe Ser Asn Tyr Ile Gly Pro Gly Asn Gln Leu Gln
 1 5 10 15
 Ala Gly Asn Pro Gln Ser Val Val Asp Ala Ala Ala Arg Ile His Asp
 20 25 30
 Phe Arg Tyr Ser Glu Leu Ile Lys Leu Gly Ile Asn Pro Tyr Thr His
 35 40 45
 Trp Ser Val Ala Asp Asp Glu Leu Leu His Asn
 50 55

<210> 22

<211> 59

<212> PRT

<213> Chipmunk parvovirus

<400> 22

Ile His Leu Pro Ala Asp Arg Tyr Leu Gly Pro Gly Asn Pro Leu Glu
 1 5 10 15
 Asn Gly Pro Pro Val Asp Pro Val Asp Ala Val Ala Arg Ile His Asp

| | | | | | | | | | | | | | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|--|--|
| | | | 20 | | | | | 25 | | | | | 30 | | | | |
| Phe | Arg | Tyr | Ala | Asp | Leu | Glu | Lys | Gln | Gly | Ile | Asn | Pro | Tyr | Thr | Thr | | |
| | | 35 | | | | | 40 | | | | | 45 | | | | | |
| Tyr | Thr | Ile | Ala | Asp | Glu | Glu | Leu | Leu | Lys | Asn | | | | | | | |
| | 50 | | | | | | 55 | | | | | | | | | | |

<210> 23

<211> 59

<212> PRT

<213> Adeno-Associated Virus 2 parvovirus

<400> 23

| | | | | | | | | | | | | | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|--|--|
| Leu | Val | Leu | Pro | Gly | Tyr | Lys | Tyr | Leu | Gly | Pro | Phe | Asn | Gly | Leu | Asp | | |
| 1 | | | | 5 | | | | | 10 | | | | | 15 | | | |
| Lys | Gly | Glu | Pro | Val | Asn | Glu | Ala | Asp | Ala | Ala | Ala | Leu | Glu | His | Asp | | |
| | | | 20 | | | | | 25 | | | | | 30 | | | | |
| Lys | Ala | Tyr | Asp | Arg | Gln | Leu | Asp | Ser | Gly | Asp | Asn | Pro | Tyr | Leu | Lys | | |
| | | 35 | | | | | 40 | | | | | 45 | | | | | |
| Tyr | Asn | His | Ala | Asp | Ala | Glu | Phe | Gln | Glu | Arg | | | | | | | |
| | 50 | | | | | | 55 | | | | | | | | | | |

<210> 24

<211> 59

<212> PRT

<213> Adeno-Associated Virus 3B parvovirus

<400> 24

| | | | | | | | | | | | | | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|--|--|
| Leu | Val | Leu | Pro | Gly | Tyr | Lys | Tyr | Leu | Gly | Pro | Gly | Asn | Gly | Leu | Asp | | |
| 1 | | | | 5 | | | | | 10 | | | | | 15 | | | |
| Lys | Gly | Glu | Pro | Val | Asn | Glu | Ala | Asp | Ala | Ala | Ala | Leu | Glu | His | Asp | | |
| | | | 20 | | | | | 25 | | | | | 30 | | | | |
| Lys | Ala | Tyr | Asp | Gln | Gln | Leu | Lys | Ala | Gly | Asp | Asn | Pro | Tyr | Leu | Lys | | |
| | | 35 | | | | | 40 | | | | | 45 | | | | | |
| Tyr | Asn | His | Ala | Asp | Ala | Glu | Phe | Gln | Glu | Arg | | | | | | | |
| | 50 | | | | | | 55 | | | | | | | | | | |

<210> 25

<211> 59

<212> PRT

<213> Adeno-Associated Virus 4 parvovirus

<400> 25

| | | | | | | | | | | | | | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|--|--|
| Leu | Val | Leu | Pro | Gly | Tyr | Lys | Tyr | Leu | Gly | Pro | Gly | Asn | Gly | Leu | Asp | | |
| 1 | | | | 5 | | | | | 10 | | | | | 15 | | | |
| Lys | Gly | Glu | Pro | Val | Asn | Ala | Ala | Asp | Ala | Ala | Ala | Leu | Glu | His | Asp | | |
| | | | 20 | | | | | 25 | | | | | 30 | | | | |
| Lys | Ala | Tyr | Asp | Gln | Gln | Leu | Lys | Ala | Gly | Asp | Asn | Pro | Tyr | Leu | Lys | | |
| | | 35 | | | | | 40 | | | | | 45 | | | | | |
| Tyr | Asn | His | Ala | Asp | Ala | Glu | Phe | Gln | Gln | Arg | | | | | | | |

50

55

<210> 26

<211> 59

<212> PRT

<213> Adeno-Associated Virus 5 parvovirus

<400> 26

Leu Val Leu Pro Gly Tyr Asn Tyr Leu Gly Pro Gly Asn Gly Leu Asp
 1 5 10 15
 Arg Gly Glu Pro Val Asn Arg Ala Asp Glu Val Ala Arg Glu His Asp
 20 25 30
 Ile Ser Tyr Asn Glu Gln Leu Glu Ala Gly Asp Asn Pro Tyr Leu Lys
 35 40 45
 Tyr Asn His Ala Asp Ala Glu Phe Gln Glu Lys
 50 55

<210> 27

<211> 59

<212> PRT

<213> Adeno-Associated Virus 6 parvovirus

<400> 27

Leu Val Leu Pro Gly Tyr Lys Tyr Leu Gly Pro Phe Asn Gly Leu Asp
 1 5 10 15
 Lys Gly Glu Pro Val Asn Ala Ala Asp Ala Ala Ala Leu Glu His Asp
 20 25 30
 Lys Ala Tyr Asp Gln Gln Leu Lys Ala Gly Asp Asn Pro Tyr Leu Arg
 35 40 45
 Tyr Asn His Ala Asp Ala Glu Phe Gln Glu Arg
 50 55

<210> 28

<211> 59

<212> PRT

<213> LuIII parvovirus

<400> 28

Trp Val Pro Pro Gly Tyr Lys Tyr Leu Gly Pro Gly Asn Ser Leu Asn
 1 5 10 15
 Gln Gly Glu Pro Thr Asn Pro Ser Asp Ala Ala Ala Lys Glu His Asp
 20 25 30
 Glu Ala Tyr Asp Gln Tyr Ile Lys Ser Gly Lys Asn Pro Tyr Leu Tyr
 35 40 45
 Phe Ser Pro Ala Asp Gln Arg Phe Ile Asp Gln
 50 55

<210> 29

<211> 59
 <212> PRT
 <213> H1 parvovirus

<400> 29

```

Trp Val Pro Pro Gly Tyr Lys Tyr Leu Gly Pro Gly Asn Ser Leu Asp
 1          5          10          15
Gln Gly Glu Pro Thr Asn Pro Ser Asp Ala Ala Ala Lys Glu His Asp
          20          25          30
Glu Ala Tyr Asp Gln Tyr Ile Lys Ser Gly Lys Asn Pro Tyr Leu Tyr
          35          40          45
Phe Ser Pro Ala Asp Gln Arg Phe Ile Asp Gln
          50          55

```

<210> 30
 <211> 59
 <212> PRT
 <213> B19 human parvovirus

<400> 30

```

Val Gln Leu Pro Gly Thr Asn Tyr Val Gly Pro Gly Asn Glu Leu Gln
 1          5          10          15
Ala Gly Pro Pro Gln Ser Ala Val Asp Ser Ala Ala Arg Ile His Asp
          20          25          30
Phe Arg Tyr Ser Gln Leu Ala Lys Leu Gly Ile Asn Pro Tyr Thr His
          35          40          45
Trp Thr Val Ala Asp Glu Glu Leu Leu Lys Asn
          50          55

```

<210> 31
 <211> 35
 <212> PRT
 <213> Naja naja snake venom phospholipase A2 IA

<400> 31

```

Asp Phe Ala Asp Tyr Gly Cys Tyr Cys Gly Arg Gly Gly Ser Gly Thr
 1          5          10          15
Pro Val Asp Asp Leu Asp Arg Cys Cys Gln Val His Asp Asn Cys Tyr
          20          25          30
Asn Glu Ala
          35

```

<210> 32
 <211> 12
 <212> PRT
 <213> Naja naja snake venom phospholipase A2 IA

<400> 32

```

Ala Val Cys Asp Cys Asp Arg Leu Ala Ala Ile Cys

```


1 5 10

<210> 33
 <211> 35
 <212> PRT
 <213> Human pancreatic phospholipase A2 IB

<400> 33
 Glu Tyr Asn Asn Tyr Gly Cys Tyr Cys Gly Leu Gly Gly Ser Gly Thr
 1 5 10 15
 Pro Val Asp Glu Leu Asp Lys Cys Cys Gln Thr His Asp Asn Cys Tyr
 20 25 30
 Asp Gln Ala
 35

<210> 34
 <211> 12
 <212> PRT
 <213> Human pancreatic phospholipase A2 IB

<400> 34
 Phe Ile Cys Asn Cys Asp Arg Asn Ala Ala Ile Cys
 1 5 10

<210> 35
 <211> 35
 <212> PRT
 <213> Human synovial fluid phospholipase A2 IIA

<400> 35
 Ser Tyr Gly Phe Tyr Gly Cys His Cys Gly Val Gly Gly Arg Gly Ser
 1 5 10 15
 Pro Lys Asp Ala Thr Asp Arg Cys Cys Val Thr His Asp Cys Cys Tyr
 20 25 30
 Lys Arg Leu
 35

<210> 36
 <211> 12
 <212> PRT
 <213> Human synovial fluid phospholipase A2 IIA

<400> 36
 Gln Leu Cys Glu Cys Asp Lys Ala Ala Ala Thr Cys
 1 5 10

<210> 37

<211> 35

<212> PRT

<213> Gaboon viper snake venom phospholipase A2 IIB

<400> 37

| | | | | | | | | | | | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Asp | Tyr | Ile | Tyr | Tyr | Gly | Cys | Tyr | Cys | Gly | Trp | Gly | Gly | Lys | Gly | Lys |
| 1 | | | | 5 | | | | | 10 | | | | 15 | | |
| Pro | Ile | Asp | Ala | Thr | Asp | Arg | Cys | Cys | Phe | Val | His | Asp | Cys | Cys | Tyr |
| | | | 20 | | | | 25 | | | | | | 30 | | |
| Gly | Lys | Met | | | | | | | | | | | | | |
| | | 35 | | | | | | | | | | | | | |

<210> 38

<211> 12

<212> PRT

<213> Gaboon viper snake venom phospholipase A2 IIB

<400> 38

| | | | | | | | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Glu | Leu | Cys | Glu | Cys | Asp | Arg | Val | Ala | Ala | Ile | Cys |
| 1 | | | | 5 | | | | | | 10 | |

<210> 39

<211> 35

<212> PRT

<213> Rat phospholipase A2 IIC

<400> 39

| | | | | | | | | | | | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Ser | Tyr | Tyr | Gly | Tyr | Gly | Cys | Tyr | Cys | Gly | Leu | Gly | Gly | Arg | Gly | Ile |
| 1 | | | | 5 | | | | | 10 | | | | | 15 | |
| Pro | Val | Asp | Ala | Thr | Asp | Arg | Cys | Cys | Trp | Ala | His | Asp | Cys | Cys | Tyr |
| | | | 20 | | | | 25 | | | | | | 30 | | |
| His | Lys | Leu | | | | | | | | | | | | | |
| | | 35 | | | | | | | | | | | | | |

<210> 40

<211> 12

<212> PRT

<213> Rat phospholipase A2 IIC

<400> 40

| | | | | | | | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Lys | Ala | Cys | Glu | Cys | Asp | Lys | Leu | Ser | Val | Tyr | Cys |
| 1 | | | | 5 | | | | | 10 | | |

<210> 41

<211> 41

<212> PRT

<213> Bee venom phospholipase A2 III

<400> 41

```

Ile Ile Tyr Pro Gly Thr Leu Trp Cys Gly His Gly Asn Lys Ser Ser
 1           5           10           15
Gly Pro Asn Glu Leu Gly Arg Phe Lys His Thr Asp Ala Cys Cys Arg
          20           25           30
Thr His Asp Met Cys Pro Asp Val Met
        35           40

```

<210> 42

<211> 12

<212> PRT

<213> Bee venom phospholipase A2 III

<400> 42

```

Leu Ser Cys Asp Cys Asp Asp Lys Phe Tyr Asp Cys
 1           5           10

```

<210> 43

<211> 35

<212> PRT

<213> Human phospholipase A2 V

<400> 43

```

Asn Tyr Gly Phe Tyr Gly Cys Tyr Cys Gly Trp Gly Gly Arg Gly Thr
 1           5           10           15
Pro Lys Asp Gly Thr Asp Trp Cys Cys Trp Ala His Asp His Cys Tyr
          20           25           30
Gly Arg Leu
        35

```

<210> 44

<211> 12

<212> PRT

<213> Human phospholipase A2 V

<400> 44

```

Asn Leu Cys Ala Cys Asp Arg Lys Leu Val Tyr Cys
 1           5           10

```

<210> 45

<211> 35

<212> PRT

<213> Human phospholipase A2 X

<400> 45

```

Ala Tyr Met Lys Tyr Gly Cys Phe Cys Gly Leu Gly Gly His Gly Gln
 1           5           10           15
Pro Arg Asp Ala Ile Asp Trp Cys Cys His Gly His Asp Cys Cys Tyr

```

20 25 30
 Thr Arg Ala
 35

<210> 46
 <211> 12
 <212> PRT
 <213> Human phospholipase A2 X

<400> 46
 Leu Leu Cys Lys Cys Asp Gln Glu Ile Ala Asn Cys
 1 5 10

<210> 47
 <211> 27
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> PCR Primer

<400> 47
 agtgggtatc gctactaacc tacactc
 27

<210> 48
 <211> 27
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> PCR Primer

<400> 48
 gatctgtcat catccagtct tctatgc
 27

<210> 49
 <211> 172
 <212> PRT
 <213> Porcine parvovirus

<400> 49
 Ala Pro Pro Ala Lys Arg Ala Arg Gly Leu Thr Leu Pro Gly Tyr Lys
 1 5 10 15
 Tyr Leu Gly Pro Gly Asn Ser Leu Asp Gln Gly Glu Pro Thr Asn Pro
 20 25 30
 Ser Asp Ala Ala Ala Lys Glu His Asp Glu Ala Tyr Asp Lys Tyr Ile
 35 40 45
 Lys Ser Gly Lys Asn Pro Tyr Phe Tyr Phe Ser Ala Ala Asp Glu Lys

| | | | | |
|-----------------------------------------------------------------|-----|-----|-----|----|
| 50 | | 55 | | 60 |
| Phe Ile Lys Glu Thr Glu His Ala Lys Asp Tyr Gly Gly Lys Ile Gly | | | | |
| 65 | 70 | 75 | 80 | |
| His Tyr Phe Phe Arg Ala Lys Arg Ala Phe Ala Pro Lys Leu Ser Glu | | | | |
| | 85 | 90 | 95 | |
| Thr Asp Ser Pro Thr Thr Ser Gln Gln Pro Glu Val Arg Arg Ser Pro | | | | |
| | 100 | 105 | 110 | |
| Arg Lys His Pro Gly Ser Lys Pro Pro Gly Lys Arg Pro Ala Pro Arg | | | | |
| | 115 | 120 | 125 | |
| His Ile Phe Ile Asn Leu Ala Lys Lys Lys Ala Lys Gly Thr Ser Asn | | | | |
| | 130 | 135 | 140 | |
| Thr Asn Ser Asn Ser Met Ser Glu Asn Val Glu Gln His Asn Pro Ile | | | | |
| 145 | 150 | 155 | 160 | |
| Asn Ala Gly Thr Glu Leu Ser Ala Thr Gly Asn Glu | | | | |
| | 165 | 170 | | |

<210> 50

<211> 516

<212> DNA

<213> Porcine parvovirus

<400> 50

gcgcttcctg caaaaagagc aagaggacta actctaccag gatacaaata ccttggtcca
60

ggaaactcac tagaccaagg agaaccaact aatccatcag acgccgcagc aaaagaacac
120

gacgaagcct acgacaaata cataaaatct ggaaaaaatc catacttcta cttctcagca
180

gctgatgaaa aattcataaa agaaactgaa cacgcaaaag actacggagg taaaattgga
240

cattacttct tcagagcaaa gcgtgccttt gctccaaaac tctcagaaac agactcacca
300

actacatctc aacaaccaga ggtaagaaga tcgccgagaa aacacccagg gtctaaacca
360

ccaggaaaaa gacctgctcc aagacatatt ttataaaact tagctaaaaa aaaagctaaa
420

gggacatcta atacaaactc taactcaatg agtgaaaatg tggaacaaca caaccctatt
480

aatgcaggca ctgaattgtc tgcaacagga aatgaa
516

<210> 51

<211> 227

<212> PRT

<213> B19 human parvovirus

<400> 51

| |
|-----------------------------------------------------------------|
| Ser Ser Lys Glu Ser Gly Lys Trp Trp Glu Ser Asp Asp Lys Phe Ala |
| 1 5 10 15 |

| |
|-----------------------------------------------------------------|
| Lys Ala Val Tyr Gln Gln Phe Val Glu Phe Tyr Lys Lys Val Thr Gly |
| 20 25 30 |

Thr Asp Leu Glu Leu Ile Gln Ile Leu Lys Asp His Tyr Asn Ile Ser
 35 40 45
 Leu Asp Asn Pro Leu Glu Asn Pro Ser Ser Leu Phe Asp Leu Val Ala
 50 55 60
 Arg Ile Lys Asn Asn Leu Lys Asn Ser Pro Asp Leu Tyr Ser His His
 65 70 75 80
 Phe Gln Ser His Gly Gln Leu Ser Asp His Pro His Ala Leu Ser Ser
 85 90 95
 Ser Ser Ser His Ala Glu Pro Arg Gly Glu Asn Ala Val Leu Ser Ser
 100 105 110
 Glu Asp Leu His Lys Pro Gly Gln Val Ser Val Gln Leu Pro Gly Thr
 115 120 125
 Asn Tyr Val Gly Pro Gly Asn Glu Leu Gln Ala Gly Pro Pro Gln Ser
 130 135 140
 Ala Val Asp Ser Ala Ala Arg Ile His Asp Phe Arg Tyr Ser Gln Leu
 145 150 155 160
 Ala Lys Leu Gly Ile Asn Pro Tyr Thr His Trp Thr Val Ala Asp Glu
 165 170 175
 Glu Leu Leu Lys Asn Ile Lys Asn Glu Thr Gly Phe Gln Ala Gln Val
 180 185 190
 Val Lys Asp Tyr Phe Thr Leu Lys Gly Ala Ala Ala Pro Val Ala His
 195 200 205
 Phe Gln Gly Ser Leu Pro Glu Val Pro Ala Tyr Asn Ala Ser Glu Lys
 210 215 220
 Tyr Pro Ser
 225

<210> 52

<211> 678

<212> DNA

<213> B19 human parvovirus

<400> 52

agtaaagaaa gtggcaaatg gtgggaaagt gatgataaat ttgctaaagc tgtgtatcag
 60

caatttgtgg aattttataa aaagggttact ggaacagact tagagcttat tcaaatatta
 120

aaagatcatt ataatatctt tttagataat cccctagaaa acccatcctc tctgtttgac
 180

ttagttgctc gtattaaaaa taaccttaaa aactctccag acttatatag tcatcatttt
 240

caaagtcattg gacagttatc tgaccacccc catgccttat catccagtag cagtcattgca
 300

gaacctagag gagaaaaatgc agtattatct agtgaagact tacacaagcc tgggcaagtt
 360

agcgtacaac taccggtac taactatggt gggcctggca atgagctaca agctgggccc
 420

ccgcaaagtg ctgttgacag tgctgcaagg attcatgact ttaggtatag ccaactggct
 480

aagttgggaa taaatccata tactcattgg actgtagcag atgaagagct tttaaaaaat
 540

ataaaaaaatg aaactggggtt tcaagcaciaa gtagtaaaaag actacttttac tttaaaaggt
 600
 gcagctgccc ctgtggccca ttttcaagga agtttgccgg aagttcccgcc ttacaacgcc
 660
 tcagaaaaaat acccaagc
 678

<210> 53

<211> 276

<212> PRT

<213> Galleria mellonella densovirus

<400> 53

| | | | | | | | | | | | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Met | Ser | Phe | Phe | Lys | Asn | Gln | Leu | Ile | His | Arg | Ala | Arg | Pro | Gly | Tyr |
| 1 | | | | 5 | | | | | 10 | | | | | 15 | |
| Arg | Ile | Ile | Pro | Glu | Ser | Thr | Val | Thr | Glu | Asp | Ile | Glu | Leu | Gly | Thr |
| | | | 20 | | | | | 25 | | | | | 30 | | |
| Ile | Gly | Glu | Glu | Thr | Pro | Leu | Leu | Ser | Glu | Gly | Val | Ile | Thr | Ala | Val |
| | | 35 | | | | | 40 | | | | | 45 | | | |
| Glu | Glu | Gly | Ala | Ile | Gly | Leu | Pro | Glu | Val | Ala | Ile | Gly | Val | Ala | Gly |
| | 50 | | | | | 55 | | | | | 60 | | | | |
| Ala | Ile | Gly | Thr | His | Ala | His | Glu | Trp | Trp | Arg | Asp | Arg | Tyr | Ala | Phe |
| 65 | | | | | 70 | | | | | 75 | | | | | 80 |
| Lys | Ser | Val | Leu | Thr | Gly | Asn | Tyr | Thr | Asp | Leu | Lys | Gly | Asn | Pro | Leu |
| | | | | 85 | | | | | 90 | | | | | 95 | |
| Lys | Pro | Arg | Asn | Ala | Ile | Pro | Glu | Lys | Ile | Lys | Gln | Leu | Gly | Lys | Lys |
| | | | 100 | | | | | 105 | | | | | 110 | | |
| Ile | Phe | Gln | Gly | Asp | Phe | Asn | Arg | Ala | Phe | Pro | Asp | Asn | Leu | Lys | Leu |
| | | 115 | | | | | 120 | | | | | 125 | | | |
| Glu | Thr | Glu | Lys | Glu | Lys | Ala | Asp | Leu | Leu | Arg | Tyr | Tyr | Asn | His | Asn |
| | 130 | | | | | 135 | | | | | 140 | | | | |
| Arg | Arg | Leu | Ala | Gly | Leu | Ser | Glu | Ala | Tyr | Pro | Gln | Gly | Lys | Gly | Tyr |
| 145 | | | | | 150 | | | | | 155 | | | | | 160 |
| Ala | Tyr | Ala | Lys | Ser | Gln | Lys | Val | Leu | Glu | Ala | Glu | Arg | Arg | Gly | Leu |
| | | | | 165 | | | | | 170 | | | | | 175 | |
| Thr | Val | Pro | Gly | Tyr | Lys | Tyr | Leu | Gly | Pro | Gly | Asn | Ser | Leu | Asn | Arg |
| | | | 180 | | | | | 185 | | | | | 190 | | |
| Gly | Gln | Pro | Ile | Asn | Gln | Ile | Asp | Glu | Asp | Ala | Lys | Glu | His | Asp | Glu |
| | | 195 | | | | | 200 | | | | | 205 | | | |
| Ala | Tyr | Asp | Lys | Val | Lys | Thr | Ser | Gln | Glu | Val | Ser | Arg | Ala | Asp | Asn |
| | 210 | | | | | 215 | | | | | 220 | | | | |
| Thr | Phe | Val | Asn | Lys | Ala | Leu | Asp | His | Val | Val | Asn | Ala | Ile | Asn | Phe |
| 225 | | | | | 230 | | | | | 235 | | | | | 240 |
| Lys | Glu | Thr | Pro | Gly | Asn | Ala | Phe | Gly | Ala | Ala | Ile | Gly | Ala | Ile | Gly |
| | | | | 245 | | | | | 250 | | | | | 255 | |
| Ile | Gly | Thr | Lys | Gln | Ala | Ile | Glu | Lys | Tyr | Ser | Gly | Val | Ile | Tyr | Pro |
| | | | 260 | | | | | 265 | | | | | 270 | | |
| Ser | Val | Ser | Gly | | | | | | | | | | | | |
| | | | 275 | | | | | | | | | | | | |

<210> 54

<211> 828
 <212> DNA
 <213> Galleria mellonella densovirus

<400> 54
 atgtctttct tcaaaaatca gttgatacat cgcgcacgac ctggttatcg tataataccg
 60
 gaaagtactg ttactgaaga tattgaatta ggtactattg gtgaagaaac tccattgtta
 120
 agtgaagggtg ttattacagc tgtagaagaa ggtgctattg gattaccaga agttgctatt
 180
 ggtgtggctg gtgctattgg aacacatgca catgaatggt ggagagatag atacgctttt
 240
 aaatctgttt taactggtaa ttatactgat ttaaaaggaa atcctttaa acctagaaac
 300
 gctattcctg aaaaaattaa acaactagga aagaaaatat ttcaaggaga ttttaatcgt
 360
 gcatttcctg ataattttaa attggaaact gaaaaagaaa aagctgattt attaagatat
 420
 tataatcata atagaagatt agctggacta agtgaagctt atccacaagg gaaaggatat
 480
 gcttatgcta aaagtcaaaa agtattagaa gctgaacgac gtggattaac tgttcctgga
 540
 tataaatatc ttggctcctgg aaattcattg aatagaggtc aacctataaa tcaaatagac
 600
 gaagacgcta aagaacacga cgaagcttat gataaagtga aaacaagtca agaagtaagt
 660
 cgagcagata atacatttgt taataaagcg ttagatcacg tggttaatgc tattaatttt
 720
 aaagaaacac ctggtaacgc ttttggtgct gctattggag ctattggaat tggaactaag
 780
 caagctattg aaaaatatag tggagtaatc tacccttctg tttcaggt
 828

<210> 55
 <211> 94
 <212> PRT
 <213> Casphalia extranea densovirus

<400> 55
 Met Pro Arg Ile His Phe Pro Tyr His Asn Tyr Leu Gly Pro Gly Ser
 1 5 10 15
 Asp Asn Phe Lys Lys Gln Pro Val Asp Glu Asp Asp Ala Ile Ala Arg
 20 25 30
 Ala His Asp Leu Asp Tyr Asp Lys Ala Ser Ser Asp Lys Asp Ile Phe
 35 40 45
 Lys Ala Asp Lys Gln Ala Arg Asp Glu Phe Phe Ser Ser Phe Val His
 50 55 60
 Ser Gly Asn Leu His Ser Leu Ile Gly Gly Leu Gly Leu Gly Ala Lys
 65 70 75 80
 Asn Leu Val Glu Glu His Val Leu Gly Lys Ser Leu Tyr Gly
 85 90

<210> 56
 <211> 282
 <212> DNA
 <213> Casphalia extranea densovirus

<400> 56
 atgcctcgta ttcatTTTcc ttatcataat tatcttggtc cgggcagtga taactTTTaaa
 60
 aaacaaccag tagacgaaga cgacgcaata gccagagcac atgacctgga ttacgataaa
 120
 gcaagctctg ataaagacat tttcaaggct gataagcagg ctcgcgacga gTTTTTcagt
 180
 tcatttTgtgc acagcggaaa cttgcatagt ttaattggTg gactaggact tggagctaaa
 240
 aatctagtag aagagcatgt actaggtAag tccttTgtacg gt
 282

<210> 57
 <211> 94
 <212> PRT
 <213> Bombyx mori densovirus

<400> 57
 Met Pro Arg Ile His Phe Pro Tyr His Asn Tyr Leu Gly Pro Gly Thr
 1 5 10 15
 Asp Asn Phe Glu Lys Asn Pro Ile Asp Glu Asp Asp Ala Ile Ala Arg
 20 25 30
 Ser His Asp Leu Ala Tyr Asp Lys Val Thr Asn His Lys Glu Val Phe
 35 40 45
 Gln Ala Asp Lys Gln Ala Arg Asp Glu Phe Phe Thr Ser Phe Val His
 50 55 60
 Thr Gly Asn Val His Ser Leu Ile Gly Gly Ile Gly Leu Gly Thr Lys
 65 70 75 80
 Asn Leu Val Glu Glu His Val Leu Gly Lys Pro Leu Tyr Gly
 85 90

<210> 58
 <211> 282
 <212> DNA
 <213> Bombyx mori densovirus

<400> 58
 atgcctcgta ttcatTTTcc ttatcataat tatcttggtc cgggtactga taactTTtgaa
 60
 aaaaatccaa tagacgaaga cgacgctatc gcgaggTcgC atgattTggc ttacgataaa
 120
 gtaaccaatc ataaggaagt ttttcaagct gataaacagg cccgtgacga gTTTTTTact
 180
 tcatttTgtgc atactggaaa cgtgcatagt ttaattggcg gtattggact tggaaactaaa
 240

aatttggtag aagaacatgt actaggtaaa cccttgtag ga
282

<210> 59

<211> 202

<212> PRT

<213> Adeno-associated virus

<400> 59

| | | | | | | | | | | | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Met | Ala | Ala | Asp | Gly | Tyr | Leu | Pro | Asp | Trp | Leu | Glu | Asp | Thr | Leu | Ser |
| 1 | | | | 5 | | | | | 10 | | | | | 15 | |
| Glu | Gly | Ile | Arg | Gln | Trp | Trp | Lys | Leu | Lys | Pro | Gly | Pro | Pro | Pro | Pro |
| | | | 20 | | | | | 25 | | | | | 30 | | |
| Lys | Pro | Ala | Glu | Arg | His | Lys | Asp | Ser | Arg | Gly | Leu | Val | Leu | Pro | |
| | | 35 | | | | | 40 | | | | 45 | | | | |
| Gly | Tyr | Lys | Tyr | Leu | Gly | Pro | Phe | Asn | Gly | Leu | Asp | Lys | Gly | Glu | Pro |
| | 50 | | | | | 55 | | | | | 60 | | | | |
| Val | Asn | Glu | Ala | Asp | Ala | Ala | Ala | Leu | Glu | His | Asp | Lys | Ala | Tyr | Asp |
| 65 | | | | | 70 | | | | | 75 | | | | | 80 |
| Arg | Gln | Leu | Asp | Ser | Gly | Asp | Asn | Pro | Tyr | Leu | Lys | Tyr | Asn | His | Ala |
| | | | | 85 | | | | | 90 | | | | | 95 | |
| Asp | Ala | Glu | Phe | Gln | Glu | Arg | Leu | Lys | Glu | Asp | Thr | Ser | Phe | Gly | Gly |
| | | | 100 | | | | | 105 | | | | | 110 | | |
| Asn | Leu | Gly | Arg | Ala | Val | Phe | Gln | Ala | Lys | Lys | Arg | Val | Leu | Glu | Pro |
| | | 115 | | | | | 120 | | | | | 125 | | | |
| Leu | Gly | Leu | Val | Glu | Glu | Pro | Val | Lys | Thr | Ala | Pro | Gly | Lys | Lys | Arg |
| | 130 | | | | | | 135 | | | | 140 | | | | |
| Pro | Val | Glu | His | Ser | Pro | Val | Glu | Pro | Asp | Ser | Ser | Ser | Gly | Thr | Gly |
| 145 | | | | | 150 | | | | | 155 | | | | | 160 |
| Lys | Ala | Gly | Gln | Gln | Pro | Ala | Arg | Lys | Arg | Leu | Asn | Phe | Gly | Gln | Thr |
| | | | 165 | | | | | | 170 | | | | | 175 | |
| Gly | Asp | Ala | Asp | Ser | Val | Pro | Asp | Pro | Gln | Pro | Leu | Gly | Gln | Pro | Pro |
| | | | 180 | | | | | 185 | | | | | 190 | | |
| Ala | Ala | Pro | Ser | Gly | Leu | Gly | Thr | Asn | Thr | | | | | | |
| | | 195 | | | | | 200 | | | | | | | | |

<210> 60

<211> 606

<212> DNA

<213> Adeno-associated virus

<400> 60

atggctgccc atggttatct tccagattgg ctcgaggaca ctctctctga aggaataaga
60
cagtgggtgga agctcaaacc tggcccacca ccaccaaagc ccgcagagcg gcataaggac
120
gacagcaggg gtcttggtgct tctgggttac aagtacctcg gacccttcaa cggactcgac
180
aaggagagc cggtcaacga ggcagacgcc gcggccctcg agcacgacaa agcctacgac
240
cggcagctcg acagcggaga caaccgtac ctcaagtaca accacgccga cgcgagagttt

300
caggagcgcc ttaaagaaga tacgtctttt gggggcaacc tcggacgagc agtcttccag
360
gcgaaaaaga gggttcttga acctctgggc ctggttgagg aacctgttaa gacggctccg
420
ggaaaaaaga ggccggtaga gcactctcct gtggagccag actcctcctc gggaaccgga
480
aaggcggggcc agcagcctgc aagaaaaaga ttgaattttg gtcagactgg agacgcagac
540
tcagtacctg acccccagcc tctcggacag ccaccagcag cccctcttgg tctgggaact
600
aatacg
606

<210> 61

<211> 276

<212> PRT

<213> Mythimna loreyi densovirus

<400> 61

| | | | | | | | | | | | | | | | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Met | Ser | Phe | Tyr | Thr | Ser | Gly | Leu | Thr | His | Arg | Ala | Arg | Pro | Gly | Tyr | 1 | 5 | 10 | 15 |
| Arg | Ile | Ile | Pro | Glu | Ser | Thr | Ala | Thr | Glu | Asp | Ile | Glu | Leu | Gly | Thr | 20 | 25 | 30 | |
| Ile | Gly | Glu | Glu | Thr | Pro | Leu | Leu | Ser | Glu | Ser | Ala | Ile | Thr | Ala | Val | 35 | 40 | 45 | |
| Glu | Glu | Gly | Ala | Ile | Gly | Val | Pro | Glu | Val | Ala | Val | Gly | Leu | Ala | Gly | 50 | 55 | 60 | |
| Ala | Ile | Gly | Thr | His | Ala | Asp | Val | Leu | Tyr | Arg | Asn | Arg | Asn | Val | Phe | 65 | 70 | 75 | 80 |
| Lys | Ser | Val | Leu | Thr | Gly | Asn | Tyr | Thr | Asp | Leu | Lys | Gly | Asn | Pro | Ile | 85 | 90 | 95 | |
| Lys | Gln | Arg | Asn | Ser | Ile | Pro | Glu | Lys | Thr | Lys | Gln | Leu | Gly | Lys | Gly | 100 | 105 | 110 | |
| Ile | Phe | Gln | Gly | Asp | Phe | Asn | Arg | Ala | Phe | Pro | Glu | Asp | Leu | Lys | Thr | 115 | 120 | 125 | |
| Glu | Thr | Glu | Gln | Glu | Lys | Asn | Asp | Leu | Leu | Arg | Tyr | Tyr | Asn | His | Asn | 130 | 135 | 140 | |
| Arg | Arg | Leu | Ala | Gly | Leu | Ser | Glu | Ala | Tyr | Pro | Gln | Gly | Lys | Gly | Tyr | 145 | 150 | 155 | 160 |
| Ala | Tyr | Ala | Lys | Ser | Gln | Lys | Val | Leu | Glu | Ala | Glu | Arg | Arg | Gly | Leu | 165 | 170 | 175 | |
| Thr | Val | Pro | Gly | Tyr | Lys | Tyr | Leu | Gly | Pro | Gly | Asn | Ser | Leu | Asn | Arg | 180 | 185 | 190 | |
| Gly | Gln | Pro | Thr | Asn | Gln | Ile | Asp | Glu | Asp | Ala | Lys | Glu | His | Asp | Glu | 195 | 200 | 205 | |
| Ala | Tyr | Asp | Lys | Ala | Lys | Thr | Ser | Gln | Glu | Val | Ser | Glu | Ala | Asp | Asn | 210 | 215 | 220 | |
| Thr | Phe | Val | Asn | Lys | Ala | Leu | Asp | His | Val | Val | Asn | Ala | Ile | Asn | Phe | 225 | 230 | 235 | 240 |
| Lys | Glu | Thr | Pro | Gly | Asn | Ala | Phe | Gly | Ala | Ala | Ile | Gly | Ala | Ile | Gly | 245 | 250 | 255 | |

Ile Gly Thr Lys Gln Ala Ile Glu Lys His Ser Gly Val Ile Tyr Pro
 260 265 270
 Ser Val Ser Gly
 275

<210> 62
 <211> 828
 <212> DNA
 <213> Mythimna loreyi densovirus

<400> 62
 atgtctttct acacgagcgg gttaacacat cgcgcgcgac ctggctatcg tataatacca
 60
 gaaagtactg ctaccgaaga tatagaatta ggtactattg gtgaagaaac tcctttatta
 120
 agtgaaagtg ctattactgc tgtagaagaa ggtgctattg gagttccaga agttgccggt
 180
 ggacttgctg gtgctatagg aacacatgct gacgtgttat atagaaatag aaacgtatct
 240
 aaaagtgttt taactggaaa ttacactgat ttaaaaggaa atccaataaa acaaagaaat
 300
 tctattcctg agaagacaaa gcaattagga aaaggtatat ttcaagggtga ttttaatcgt
 360
 gcatttcctg aagattttaa aacagaaact gagcaagaaa aaaacgattt attacgatac
 420
 tataatcata atagaagact agctggttta agtgaagctt atccacaagg aaaaggatac
 480
 gcttatgcta agagtcaaaa agtattagaa gctgaaagac gtggattaac tgttcctgga
 540
 tataaatatc ttggctcctg aaactcactt aatagaggtc aacctactaa tcaaatagac
 600
 gaagacgcta aagaacacga cgaagcttac gataaagcaa aaacaagtca agaagtaagt
 660
 gaagcagata atacatttgt taataaggcg ttagatcacg tggttaatgc tatcaatttt
 720
 aaagaaacgc ctggtaacgc ttttggtgct gctattggag ctattggaat tggaactaag
 780
 caagctattg aaaaacacag tggagtaatc tacccttctg tttcaggt
 828

<210> 63
 <211> 268
 <212> PRT
 <213> Acheta domesticus densovirus

<400> 63
 Met Ser Gly Val Phe Thr Asp Leu Thr Leu Arg Asp Phe Ser Leu Gly
 1 5 10 15
 Thr Asn Asn Leu Pro Leu Ala Asn Thr Pro Lys Leu Arg Asn Arg Phe
 20 25 30
 Gly Arg Trp Trp Asn Arg Ser His Pro Tyr Asp Arg Leu Pro Thr Asn

| | | | | | | | | | | | | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|--|
| | | 35 | | | | | 40 | | | | | 45 | | | | |
| Glu | Pro | Glu | Pro | Leu | Arg | Glu | Thr | Ser | Phe | Gln | Glu | Ala | Ala | Gly | Pro | |
| | 50 | | | | | 55 | | | | | 60 | | | | | |
| Glu | Glu | Thr | Arg | Ile | Asp | Ile | Ala | Glu | Asp | Glu | Ile | Asn | Ala | Gly | Glu | |
| 65 | | | | | 70 | | | | | 75 | | | | | 80 | |
| Gly | Ala | Ala | Glu | Ala | Glu | Thr | Ser | Phe | Ser | Thr | Gly | Val | Glu | Glu | Thr | |
| | | | | 85 | | | | | 90 | | | | | 95 | | |
| Ala | Leu | Glu | Leu | Gly | Glu | Ala | Ala | Thr | Glu | Thr | Thr | Gly | Leu | Leu | Gly | |
| | | | 100 | | | | | 105 | | | | | 110 | | | |
| Gly | Ala | Thr | Ala | Gly | Ser | Ala | Ala | Ala | Gly | Thr | Ala | Gly | Thr | Leu | Gly | |
| | | 115 | | | | | | 120 | | | | 125 | | | | |
| Thr | Val | Ala | Ala | Thr | Ala | Ala | Gly | Gly | Ala | Ala | Leu | Ala | Gly | Ile | Gly | |
| | 130 | | | | | 135 | | | | | 140 | | | | | |
| Ile | Gly | Ile | Lys | Lys | Leu | Ile | Asp | His | Thr | Ser | Ser | Lys | Gly | Ala | Val | |
| 145 | | | | | 150 | | | | | 155 | | | | | 160 | |
| Leu | Pro | Gly | Thr | Asp | Phe | Val | Gly | Pro | Gly | Asn | Pro | Ile | Asp | Pro | Lys | |
| | | | | 165 | | | | | 170 | | | | | 175 | | |
| Pro | Ala | Arg | Ser | Glu | Thr | Asp | Gln | Ile | Ala | Lys | Glu | His | Asp | Leu | Gly | |
| | | | 180 | | | | | 185 | | | | | 190 | | | |
| Tyr | Glu | Asp | Leu | Leu | His | Arg | Ala | Lys | Ser | Gln | Tyr | Phe | Thr | Glu | Glu | |
| | | 195 | | | | | 200 | | | | | 205 | | | | |
| Asp | Phe | Lys | Thr | Glu | Val | Tyr | Lys | Leu | Asp | Asp | Glu | Ala | Ile | His | Arg | |
| | 210 | | | | | 215 | | | | | 220 | | | | | |
| Phe | Ser | Glu | Glu | Tyr | Gln | Lys | Ser | Gly | Thr | Trp | Gln | Ala | Phe | Val | Gly | |
| 225 | | | | | 230 | | | | | 235 | | | | | 240 | |
| Lys | Tyr | Gly | Leu | Lys | Ala | Lys | Arg | Val | Ile | Glu | Asp | Val | Ile | Gly | Gly | |
| | | | | 245 | | | | | 250 | | | | | 255 | | |
| Pro | Val | Tyr | Pro | Gln | Gln | Pro | Lys | Lys | Arg | Glu | Tyr | | | | | |
| | | | 260 | | | | | 265 | | | | | | | | |

<210> 64

<211> 804

<212> DNA

<213> Acheta domesticus densovirus

<400> 64

atgtctggcg tctttacaga tctcacgttg cgtgatttca gtttaggtac taataattta
60

cctttagcaa atacacctaa acttcgtaat agattttggga gatggtggaa tcgtagtcac
120

ccctacgaca gactacctac aaatgaacca gaacctcttc gcgaaactag tttccaagag
180

gcggcagggc cagaagaaac tcgcattgat atcgcggaag acgaaatcaa cgcagggtgaa
240

ggagcagccg aagctgaaac aagtttctct actggagttg aggaaacagc acttgaactc
300

ggcgaagcag ctaccgaaac aacaggactt ttgggaggcg ctactgcagg .tagcgcagca
360

gcaggcacag ctggaactct tggaactgtc gcggcaactg cagcaggagg agcagccctc
420

gcaggaatcg qcatcgaat taaaaaatta attgatcaca cgagtaqcaa aggtgcagta

480
 ttacctggca ctgatttcgt ggggcctggt aatccaatag atcctaaacc tgcacgatca
 540
 gaaactgata aaattgctaa ggaacacgat cttgggtacg aagacttatt gcatcgtgca
 600
 aaatcacagt attttactga agaagatttc aaaactgaag tatataaatt ggacgacgaa
 660
 gcaattcatc gtttttctga ggaataccag aaatccggtg cgtggcaagc atttggtgga
 720
 aaatacggac ttaaagctaa acgtgtaatt gaagacgtaa ttggtggacc tgtatatcct
 780
 caacaaccta aaaaacgtga gtac
 804

<210> 65

<211> 5234

<212> DNA

<213> *Acheta domesticus* densovirus

<400> 65

ttatatcctt aggatataaa gagcaagcac ccaggcactt cttctattct gcctcagatc
 60
 gtcgatataa gcggagcttt tctgtgactg tacgtagtgt attgatcttc gtgttctgga
 120
 actcaggtat gttgttcgct gcagtagaag cgcctccagc atgtacgttc gaacctcgtt
 180
 cattgcaaca gttatcactt agaacttttg tagaaaatac ggattgggac gtaataaatt
 240
 ttctctataa aaataatgaa ttttctagca gggtgtattc atcgatgttt tatgaaagaa
 300
 aatatctgga taatccgcat gaatatcatg attggtcagt accaaataat gttcccattg
 360
 aaacgtacat aaagataatg aattatcaca tggaagacgg tattccatca tttgctttcc
 420
 aacgcaattt agtgaataga tgtttcatac aagtgcattt ctgtgctcct ggtacttttg
 480
 gttccctat tacagataat atttgtatac cttgttataa tacatttgcg aggtttaaca
 540
 gctctaattt ctatcggttt aagaagtttg ttctcgttcg cgatcaaaca gttattgaag
 600
 atgatgaact tatcgattat atgcaatgta gaagtaattg gtgtacttgt tgtaatacaa
 660
 catcattatt tgcgattaaa gatggaactc atttaggccc gcagtggttt aatagtatat
 720
 atctcatgtg taacgaaatt ccgttattac ctgttgacga tgataactga gttaaacgat
 780
 ggagccagta caagcacagg acaatctgga tggagtagag gagacgttga aggactttat
 840
 gaagacatta tggggcctct gcgagaatca accgttgacc aatccgcaat ggtgggcgga
 900
 catgctggag agtgttacca tttctctgga ggatcagaac acgaacctag cctcagatta
 960

cgcacaaatta aagaaaagct tcctgaagat tatatcgac tggcagacaa tgtcaacgaa
1020
atggttgggc acttcattcg cggacttaga tcgaagagtt tcagctctac tcggcggtac
1080
atttcagacg ttgtggtcac gtcgtctcct gaacacgggc gcagagtgcg ggacttcctt
1140
aataagtgcg caagcactta ccccggaaga ctactactct gggtaacga aggagaccac
1200
atccacgtcg tccacgactg cccgtacagc gctggacaat gtagatgcaa gttctcaaaa
1260
acggaagact tcaaaaaggg acttcgcaag cctctacgca agtcgaagtt cattaccgaa
1320
ctcgatgaca ccgactgggc aaatgttttc ttatacttca ttgtgtcaaa atggaaaagc
1380
gaacgagcag tttggattga tggagcctta cgacgactac cggatacagg tgaagggtga
1440
cgatggaccg cgctgtgcga gcaatccga ctactactgg aacaacaagc tgcaagagat
1500
ggatgtgacg gtatcgaaga agtcacagat ttggaaggac atcaacaacg cgttttgcgt
1560
ggcgataacc aacctcgaag aaaacggagc aagttccaga ctttcgccga agaagcaaaa
1620
gtactattct caaaattgaa ctgtattcct gtgcgtgacg ctagaataat aattgatagg
1680
aagcatccta tgtattttga attgcatgat ccaaaaaatg aaaagttgta tactgctgct
1740
atttctttat atgtgcgtga cattaatgaa atgtctttga atgatttcta cttacagact
1800
ttagagaata catgtgttta tcaagctggc aatatagacc cttttgtttt ttatcataat
1860
atggaagata gtgttcactt tcttaatgaa cttatcttat ttcaatgtaa tggtagtgag
1920
gaagggtgta aagaattact aatgaatatt gtagcttggc ttaatagggt aggttggtat
1980
atgccaacaa ttaatatgaa tggggaaaaa gaatatcagc ttaaccctaa agttaatact
2040
gtatgtataa ttggtaatca taattgtggc aagaattatt tttgggacgc agtttgctgt
2100
ttgggtttta atgtggggtt tcttggtaga gttaataata aaactaacia gtttgctttg
2160
caagattgtg ttcacaaacg aattgtaatt ggtaatgaaa tatctttgga agatggagca
2220
aaggaggata tgaagaaatt atgtgagggg tgtccattca atgttagcgt aaaacatcaa
2280
tcagatggaa ttgttgccag gacacctgtt tgcttaatta gtaacaattg tatacctatt
2340
gcatgtgatg ctcatcttgc agatgtaagg atgaaggat tttattggag gccatgtagt
2400
tatttggcca aatcaaggaa gaagccgtat ccaccggctg tttttgaatt aatgaaaatg
2460
tatggagttg attggaataa agtttattaa ttccttttat tacatctttc atttttgttg
2520

caatcccata atagtactct tcccaactgc atgagtgtac tccaaagtag aactggaata
2580
aatctgggta tcaggaatag tattaggacc gttaaaaaga ggtctaattg aaggatattg
2640
aagtcgtact ttgcaagcag cttcaacaat aaagtaagct tgagtgtcag taaaattagc
2700
attattaaga ttatcatcta aattagctgt tgtcaatgca tgaacagggt tcacaccaat
2760
atgcaaacta gggtgagctt ttgggtgtata agatggaaaa gctcctgcat gaagtatttg
2820
acttttttca atttgagtcg caatagtcca ttcagtattt gatactacag tatcattaaa
2880
tttttctgaa tcccctctta aaatattatt agtcatagtg aaatcagcag ttcgatactg
2940
cgagttccct gtacctgctg ggcaactaag agttgagtta ccaccactg cgctagggtac
3000
accagtatat acagctggta tagctttttt aattaaacct aattgtgggt tataattata
3060
ttcaacaatt tctcctggag gtccattaca tctaatttca tttacatgag attgtaaaca
3120
ttcgtaacct ggattatctg ccgtaccgtt ctggtgaaga acaagtgcga aataattctt
3180
taatggatac ggaataccga actgatgacg tgggtacatga ttgggtactg atttatcgcc
3240
gtaaaactct ttaacataat cctcaaaaag attttcatca cgtattgctc caagttcatc
3300
tatggatgat acaaccattg gttcattagc ttgaaattct ttaggacgta cgtcaatacc
3360
ttgtgttttt atattcaggc ctgtagcatg aagaataaat tgattttgat ttagagtagc
3420
aagaccagta ttagaactat ttgtttgaaa agcaatacga ggggtaaatg ctgttacgcg
3480
tacattacat tcttcaatgc aacttccagg tggtaattga gcaaattcat ttggattcat
3540
atacaagaac ggtcggtcga ctggtaaacy tcccaaacta gtaataacga agttgtcaac
3600
atatgtaaca tcgttaactg tccttgttat tgggaattaca acatttgcaa gaccataagt
3660
taaaaatcga tgtactttgt taaagataat gaaattgtta tgcggagtag taattggtct
3720
aggaattgga ataatcgctt gaccgcctcc ttcacctgca atatctgcac ctgctcccg
3780
taatgatgta cctgccattc ttgaacgttt accaccacta cttttaggag gacgaccacc
3840
accgccgctt ttgccctttc ctttgtcttg tgttttgtcc ggggttatgg gaggagttaa
3900
aggagcactc atggacgaag aatcaccttc gtcagggtga gtactatcta caggattact
3960
tacaggttca ggatctggat ctgtggaacg tccagggtgc gctgctaaat cagtagtcgg
4020
atgatgttcc tgtgtagtag aaggaagttg atctacgtcg aagtcgtcaa tgtcttccgg
4080

attaccttca ggtagctcgt ctaatatatc atctgcagtt ggagttgcag gtgcgacttc
4140
ttcttcttct gcacgaggag gttctggaat aggatgatca attggtaaac ctgcacgtac
4200
tcgagctaatt tgccattggt caacagcgta ccgccggtga ccttcattta agtattccca
4260
attatcacgt tcttgtggat gaggtttcag tgggggtgtc attacactga aaacaaaata
4320
aaaagtatca gtactcacgt tttttaggtt gttgaggata tacaggcca ccaattacgt
4380
cttcaattac acgttttagct ttaagtccgt attttccaac aaatgcttgc cacgtaccgg
4440
atttctggta ttcctcagaa aaacgatgaa ttgcttcgtc gtccaattta tatacttcag
4500
ttttgaaatc ttcttcagta aaatactgtg attttgcacg atgcaataag tcttcgtacc
4560
caagatcgtg ttccttagca atttgatcag tttctgatcg tgcaggttta ggatctattg
4620
gattaccagg cccacgaaa tcagtccag gtaatactgc acctttgcta ctogtgtgat
4680
caattaattt ttttaattccg atgccgattc ctgcgagggc tgctcctcct gctgcagttg
4740
ccgcgacagt tccaagagtt ccagctgtgc ctgctgctgc gctacctgca gtagcgcctc
4800
ccaaaagtcc tgttgtttcg gtagctgctt ccgcgagttc aagtgtgtt tctcaactc
4860
cagtagagaa acttgtttca gcttcggctg ctccctcacc tgcgttgatt togtcttcg
4920
cgatatcaat gcgagtttct tctggccctg ccgcctcttg gaaactagtt togcgaagag
4980
gttctggttc atttgtaggt agtctgtcgt aggggtgact acgattccac catctcccaa
5040
atctattacg aagtttaggt gtatttgcta aaggtaaatt attagtacct aaactgaaat
5100
cacgcaacgt gagatctgta aagacgccag acatttaata ctacgatgct aaaacttccg
5160
cttatatacc ttctgacgtc agcgaaatag aagaagtgcc tgggtgcttg ctctttatat
5220
ccttcggata taaa
5234

<210> 66
<211> 36
<212> DNA
<213> unknown

<400> 66
tgggtctagtt cgcattatcc tccgcattgg actgcg
36

<210> 67
<211> 12
<212> PRT
<213> unknown

<400> 67
Trp Ser Ser Ser His Tyr Pro Pro His Trp Thr Ala
1 5 10

<210> 68
<211> 36
<212> DNA
<213> unknown

<400> 68
gtgaatcagt cttatacttc gacttggttt tgggct
36

<210> 69
<211> 12
<212> PRT
<213> unknown

<400> 69
Val Asn Gln Ser Tyr Thr Ser Thr Trp Phe Trp Ala
1 5 10

<210> 70
<211> 36
<212> DNA
<213> unknown

<400> 70
tggcctcagt tgtttacttt tccgtgttgt aatcct
36

<210> 71
<211> 12
<212> PRT
<213> unknown

<400> 71
Trp Pro Gln Leu Phe Thr Phe Pro Cys Cys Asn Pro
1 5 10

<210> 72
<211> 36
<212> DNA
<213> unknown

<400> 72

ggtctgaaga tttggagttt gccgccgcat catggg
36

<210> 73

<211> 12

<212> PRT

<213> unknown

<400> 73

Gly Leu Lys Ile Trp Ser Leu Pro Pro His His Gly
1 5 10

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
3 January 2002 (03.01.2002)

PCT

(10) International Publication Number
WO 02/00924 A3

(51) International Patent Classification⁷: C12N 9/20,
15/55, 15/11, 7/01, C07K 16/10, C12Q 1/34

(74) Agent: MBM & CO.; P.O. Box 809, Station B, Ottawa,
Ontario K1P 5P9 (CA).

(21) International Application Number: PCT/CA01/00932

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(22) International Filing Date: 28 June 2001 (28.06.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
2,310,263 28 June 2000 (28.06.2000) CA

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): INSTITUT NATIONAL DE LA RECHERCHE SCIENTIFIQUE [CA/CA]; Tour de la Cité, 2600 Boulevard Laurier, Bureau 640, Case Postale 7500, Sainte-Foy, Québec G1V 4C7 (CA).

Published:
— with international search report

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): TIJSEN, Peter [NL/CA]; 76 Winston Circle, Pointe-Claire, Québec H9S 4X6 (CA). ZADORI, Zoltan [HU/CA]; 3215 Boulevard Cartier Ouest, App. 106, Laval, Québec H7V 1J8 (CA).

(88) Date of publication of the international search report:
11 April 2002

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 02/00924 A3

(54) Title: VIRAL PHOSPHOLIPASE A₂ ENZYMES, ANTI-VIRAL AGENTS AND METHODS OF USE

(57) Abstract: The present invention provides a novel class of phospholipase A₂ enzyme (PLA₂) from viruses and a method for identifying, isolating, purifying and characterizing enzymes of this class. The present invention includes viral PLA₂ proteins, nucleic acids, and antisense oligonucleotides, and the use of these molecules in screening methods for anti-viral agents, in decreasing the infectivity and/or replication of viruses and as research tools. The present invention further includes treatment or prevention of virus-associated diseases using viral PLA₂ inhibitors and the use of the viral PLA₂-encoding region to improve virus-based gene therapy vectors.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/CA 01/00932

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N9/20 C12N15/55 C12N15/11 C12N7/01 C07K16/10
C12Q1/34

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N C07K C12Q

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

BIOSIS, EMBASE, CHEM ABS Data, SEQUENCE SEARCH

C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category * | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------|
| X | <p>BERGERON J ET AL: "Genome organization of the Kresse strain of porcine parvovirus: Identification of the allotropic determinant and comparison with those of NADL-2 and field isolates." JOURNAL OF VIROLOGY, vol. 70, no. 4, 1996, pages 2508-2515, XP002184679 ISSN: 0022-538X * see Fig. 1 * the whole document</p> <p style="text-align: center;">--- -/-</p> | 1-33 |

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *G* document member of the same patent family

Date of the actual completion of the international search

4 December 2001

Date of mailing of the international search report

18/12/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl.
Fax: (+31-70) 340-3016

Authorized officer

Hillenbrand, G

INTERNATIONAL SEARCH REPORT

International Application No.

PCI/CA 01/00932

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

| Category * | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------|
| X | <p>ERDMAN DEAN D ET AL: "Genetic diversity of human parvovirus B19: Sequence analysis of the VP1/VP2 gene from multiple isolates."</p> <p>JOURNAL OF GENERAL VIROLOGY, vol. 77, no. 11, 1996, pages 2767-2774, XP001037602</p> <p>ISSN: 0022-1317</p> <p>* see Table 1, U38546 *</p> <p>the whole document</p> <p>---</p> | 1-33 |
| X | <p>DATABASE EMBL/GENBANK/DBJ 'Online! Accession numbers L32896 and Q90125, 29 November 1994 (1994-11-29)</p> <p>TIJSEN, P.: "Organization and expression of the ambisense genome of densovirus of Galleria mellonella (GmDNV)"</p> <p>XP002184680</p> <p>abstract</p> <p>---</p> | 1-33 |
| X | <p>DATABASE EMBL/GENBANK/DBJ 'Online! Accession numbers AB042597 and Q9JFY0, 23 May 2000 (2000-05-23)</p> <p>NONAKA, K. ET AL.: "Complete nucleotide sequence and genome organization of a newly isolated Bombyx densovirus, which is clearly different in ORF structure from BmDNV previously reported."</p> <p>XP002184681</p> <p>abstract</p> <p>---</p> | 1-33 |
| X | <p>WO 00 28004 A (RABINOWITZ JOSEPH E ; SAMULSKI RICHARD JUDE (US); UNIV NORTH CAROLI) 18 May 2000 (2000-05-18)</p> <p>the whole document</p> <p>-----</p> | 1-33 |

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 36-42, 44-62

Present claims 36-42 and 44-62 relate to anti-viral agents/peptides defined by reference to a desirable characteristic or property, namely the inhibition of viral phospholipase A2.

The claims cover all agents (and their uses) having this characteristic or property, whereas the application provides support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT for only a very limited number of such products. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible.

Independent of the above reasoning, the claims also lack clarity (Article 6 PCT). An attempt is made to define the claimed agents by reference to a result to be achieved. Again, this lack of clarity in the present case is such as to render a meaningful search of the claimed matter impossible. It appears that the claims embrace a multiplicity of known agents which may exhibit such an anti-viral activity. Thus, no meaningful search could be carried out.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/CA 01/00932

| Patent document cited in search report | Publication date | Patent family member(s) | Publication date |
|-------------------------------------------|---------------------|----------------------------|---------------------|
| WO 0028004 A | 18-05-2000 | AU 1911100 A | 29-05-2000 |
| | | EP 1135468 A1 | 26-09-2001 |
| | | WO 0028004 A1 | 18-05-2000 |
| <hr/> | | | |